



Number and Calculation Policy

Reviewed: February 2022

Next Review: February 2023

Contents:

Goldsmith Primary Academy – Maths Approach on a Page	- 3
ARK Maths Mastery - Progression in Calculations Document	- 6
ARK Maths Mastery - Progression in Vocabulary Document	- 79

Intent:

At Goldsmith we develop competent mathematicians who are equipped with the appropriate knowledge and skills to succeed in the modern world. This curriculum covers all content of the National Curriculum (2014). Through this curriculum, we develop fluent mathematicians who challenge themselves by reasoning and problem solving, as a matter of course, in each lesson. We endeavour to model mathematical structures, using concrete apparatus, enabling pupils to represent their mathematical knowledge in pictorial or abstract form.

Implementation:

• **What is our curriculum:**

In our school, we have implemented the ARK Mastery Maths Programme of Study for primary pupils. This is a highly practical and visual approach to a mastery curriculum and enables pupils to visualise problems in addition to solving them. Details of the subject matter being taught can be found in the ARK Mastery Maths Overview document. Additionally, details of how we teach calculations can be found in the ARK Mastery Maths Progression in Calculations document. We can divide our curriculum into five sub-curricula:

- ARK Mastery Maths Content Teaching
- ARK Mastery Maths Meetings
- ARK Same Day Catch-Up
- Multiplication Tables
- Arithmetic

ARK Mastery Maths Content:

EYFS - As per Development Matters, pupils in Nursery and Goldilocks are encouraged to investigate numbers and quantities in engaging, hands-on activities. Pupils are encouraged to make links between numbers and associated quantities of objects or manipulatives. In Reception, pupils are moved on to the ARK curriculum for EYFS Maths, receiving two maths lessons per week which are adult led, resulting in either an adult led activity to consolidate new knowledge or independent opportunities in Enhanced Provision. Children are also able to engage in maths-led activities throughout the week, which are based on the topic being covered. On Friday, teachers in Reception engage in personalised learning, ensuring that any gaps in knowledge are addressed quickly.

KS1 and KS2 - In KS1 and 2, pupils undertake four ARK Mastery Maths lessons per week. Teaching is borne of the concrete, pictorial, abstract approach to mathematics teaching and enables pupils to demonstrate their new knowledge in a variety of ways, using manipulatives. Pupils' tasks are differentiated in order to enable ALL to achieve the learning objective. We use a simple Green (Working Towards), Yellow (Working At) and Red (Working Beyond) system, to signify the tasks being completed. Pupils are actively encouraged to move through the colour groups, should they be confident with a concept being taught. All pupils are aware they should be aiming for the yellow task or beyond. Manipulative maths equipment is used in all maths lessons, either by direct instruction of the teacher or available on the maths "Enable Station." Equipment designed for regular use is contained within a 1:2 CPA toolkit box.

ARK Mastery Maths Meetings:

KS1 and KS2 - Pupils take part in "Maths Meetings" for 15 minutes in the afternoon session (3x/week in KS1 4x/week in KS2). These sessions are deliberately set apart from the main content teaching as they focus on engaging and activating prior knowledge. In all year groups, pupils are assessed using a grid of questions. Teachers then gap analyse this information and teach the identified gaps in subsequent maths meetings. Pupils then undertake a Maths Meeting assessment every two weeks, which should illustrate that teaching has addressed knowledge gaps. Multiplication table practice will also take place in these sessions.

ARK Same Day Catch-Up:

KS1 and KS2 - Pupils who do not succeed during teaching sessions will be identified by the class teacher to receive an 'SDCU' intervention. Pupils will work with a teacher or teaching assistant to have a second chance to learn a concept, the same day it has been taught. This takes place 4x/week in KS1 and KS2.

Multiplication Tables:

EYFS - Pupils have access to a "Numbots" login which teachers may use as an activity during free-flow play or as a direct teaching tool, later in reception. Numbots enables pupils to experience numbers in different context and begin to make simple addition and subtraction sentences, as pupils progress through the "Number Story."

KS1 - Pupils are given individual access to "Times Tables Rock Stars." Pupils are able to access this at home and in school during one maths meeting per week. In school and across our academy trust, we have frequent competitions to ensure engagement.

KS2 - As in KS1, pupils have access to their own TTRS login. Pupils are able to use this in school during one of their timetabled maths meetings. In addition, pupils in KS2 take part in the weekly 99 Club. Children demonstrate progress by completing a given number of questions in a given time. Pupils receive a certificate on successful graduation from a club. KS2 pupils will undertake the 99 Club in maths meeting time.

Arithmetic:

KS1 and KS2 - Arithmetic is timetabled as the fifth lesson of the week and is used in different ways on a bi-weekly basis. In week 1, pupils will be assessed on their arithmetic knowledge bi-weekly. In the initial session, pupils will undertake the test and then peer and shared mark the assessment, with the teacher modelling answers. In week 2, children will complete their Maths Meeting Assessment (30 mins) and take part in an arithmetic gap filling session based on the gap analysis of the previous week's arithmetic test data.

Impact:

- **How is impact measured?**

ARK Mastery Maths Content, ARK Mastery Maths Meetings and Arithmetic:

Impact of our curriculum is measured in several ways. On a daily basis, pupils' progress is measured through our #FeedForward policy; pupils are given next steps verbally and in their books. At the beginning and end of the unit, pupils are given pre- and post-unit quizzes. This enables teachers to plan for pupils' needs and identify and address gaps, where necessary. On a termly basis, pupils sit NTS mathematics assessments in arithmetic and reasoning. This data is also gap analysed in order to inform future planning and maths meeting foci. Impact of teaching is also quality assured through learning walks and lesson observation. Finally, the impact of the pupil's whole mathematical journey is measured in end-of-key-stage assessments in the EYFS, KS1 and KS2. This data is published nationally.

Multiplication Tables:

Impact of our multiplication tables curriculum is measured by progress pupils make; through levels on TTRS, Clubs in the 99 club and more formally, in the Year 4 Multiplication Tables Check.

- **How do we know children have made progress?**

There are several means of identifying progress in mathematics. The simplest is to identify the progression of pupils through the coloured groups on offer in our mathematics lessons. As previously mentioned, our pupils know they are aiming for the yellow task - working at year group standard. Pupils standardised scores, derived from performance in NTS assessments, are also key in identifying areas for development. As part of the GADY project (Goldsmith Attainment in Disadvantaged Youngsters), pupils are required to improve their standardised score by a set number of points in each term of assessment. This enables staff to see the important part they are playing in any child's progression through the math curriculum.

- **How do we challenge and support lack of progress?**


Teachers are challenged and supported through pupil's lack of progress during pupil progress meetings. Teachers are able to highlight any concerns about a child, whilst providing solutions to assist that child in their continued study. SLT and the teacher together, may decide upon offering additional intervention to the pupil. Teachers and SLT will also endeavour to engage home in supporting in school learning, through home learning and IEP home

packs.

How are we inclusive?

We believe every child should be able to access our maths curriculum. We feel that by offering quality first teaching to all, in addition to targeted and appropriate use of concrete resources, pictorial representations and eventual exposure to abstract calculations, all pupils can succeed.

In all classrooms, CPA toolkits can be found which allows children 1:2 access to the equipment used in many mathematical teaching models (Numicon, Base 10, Place Value Counters, Bead Strings)



Should pupils struggle in mathematics, additional support will always be given by teachers and teaching assistants in order to ensure that pupils achieve. Pupils are targeted for same day intervention, 4x/week, to be given a second opportunity to grasp a concept. If necessary, pupils will be identified as part of pupil progress meeting procedures and will be given extra intervention.



Progression in calculations

Year 1 – Year 6*

NB. Users should familiarise themselves with the introduction (pp 2-10) to this document before referring to individual year group guidance.

**Progression guidance is not provided for EYFS/Reception since the focus should be on the understanding of early number concepts and number sense through the use of concrete manipulatives, as exemplified in the programmes of study.*

Introduction

At the centre of the mastery approach to the teaching of mathematics is the belief that all pupils have the potential to succeed. They should have access to the same curriculum content and, rather than being extended with new learning, they should deepen their conceptual understanding by tackling challenging and varied problems. Similarly, with calculation strategies, pupils must not simply rote learn procedures but demonstrate their understanding of these procedures through the use of concrete materials and pictorial representations. This document outlines the different calculation strategies that should be taught and used in Years 1 to 6, in line with the requirements of the 2014 Primary National Curriculum.

Background

The 2014 Primary National Curriculum for mathematics differs from its predecessor in many ways. Alongside the end of Key Stage year expectations, there are suggested goals for each year; there is also an emphasis on depth before breadth and a greater expectation of what pupils should achieve.

One of the key differences is the level of detail included, indicating what pupils should be learning and when. This is suggested content for each year group, but schools have been given autonomy to introduce content earlier or later, with the expectation that by the end of each key stage the required content has been covered.

For example, in Year 2, it is suggested that pupils should be able to ‘add and subtract one-digit and two-digit numbers to 20, including zero’ and a few years later, in Year 5, they should be able to ‘add and subtract whole numbers with more than four digits, including using formal written methods (columnar addition and subtraction)’.

In many ways, these specific objectives make it easier for teachers to plan a coherent approach to the development of pupils’ calculation skills, and the expectation of using formal methods is rightly coupled with the explicit requirement for pupils to use multiple representations, including concrete manipulatives and images or diagrams – a key component of the mastery approach.

Purpose

The purpose of this document is threefold. Firstly, in this introduction, it outlines the structures for calculations, which enable teachers to systematically plan problem contexts for calculations to ensure pupils are exposed to both standard and non-standard problems. Secondly, it makes teachers aware of the strategies that pupils are formally taught within each year group, which will support them to perform mental and written calculations. Finally, it supports teachers in identifying appropriate pictorial representations and concrete materials to help develop understanding.

The policy only details the strategies; teachers must plan opportunities for pupils to apply these, for example, when solving problems, or where opportunities emerge elsewhere in the curriculum.

How to use the document

For each of the four rules of number, different strategies are laid out, together with examples of what concrete materials can be used and how, along with suggested pictorial representations. Please note that the concrete and pictorial representation examples are not exhaustive, and teachers and pupils may well come up with alternatives. The purpose of using multiple representations is to give pupils a deep understanding of a mathematical concept and they should be able to work with and explain concrete, pictorial and abstract representations, and explain the links between different representations. Depth of understanding is achieved by moving between these representations. For example, if a child has started to use a pictorial representation, it does not mean that the concrete

cannot be used alongside the pictorial. If a child is working in the abstract, depth can be evidenced by asking them to exemplify their abstract working using a concrete or pictorial representation and to explain what they have done using the correct mathematical vocabulary; language is, of course, one abstract representation but is given particular significance in the national curriculum.

Mathematical language

The 2014 National Curriculum is explicit in articulating the importance of pupils using the correct mathematical language as a central part of their learning. Indeed, in certain year groups, the non-statutory guidance highlights the requirement for pupils to extend their language around certain concepts.

“The quality and variety of language that pupils hear and speak are key factors in developing their mathematical vocabulary and presenting a mathematical justification, argument or proof.”

2014 Maths Programme of Study

Suggested language structures accompany each strategy outlined in this document. These build on one another systematically, which supports pupils in making links between and across strategies as they progress through primary school.

✓	✗
ones	units
is equal to	equals / makes
zero	oh (the letter O)

New vocabulary should be introduced in a suitable context (for example, with relevant real objects, manipulatives, pictures or diagrams) and explained precisely. High expectations of the mathematical language used are essential, with teachers modelling accurate mathematical vocabulary and expecting pupils' responses to include it ***in full sentences***.

Presentation of calculations

You will see that throughout this document, calculations are presented in a variety of ways. It is important for pupils' mathematical understanding to experience and work with calculations and missing numbers in different positions relative to the = symbol. Examples used in classwork and independent work should reflect this.

Estimation

Pupils are expected to use their developing number sense from Year 1 to make predictions about the answers to their calculations. As their range of mental strategies increases, these predictions and, later, estimates should become increasingly sophisticated and accurate. All teaching of calculation should emphasise the importance of making and using these estimates to check, first, the sense and, later, the accuracy of their calculations.

Developing number sense

Fluency in arithmetic is underpinned by a good sense of number and an ability to understand numbers as both a concept (e.g. 7 is the value assigned to a set of seven objects) and as something resulting from a process (three beads and four more beads gives seven beads altogether or $3 + 4 = 7$). Understanding that a number can be partitioned in many ways (e.g. $7 = 4 + 3$; $5 + 2 = 7$; $1 + 6 = 7$) is key to being able to use numbers flexibly in calculating strategies. The part-whole model and, later, bar models, are particularly useful for developing a relational understanding of number. Pupils who are fluent in number bonds (initially within ten and then within twenty) will be able to use the 'Make ten' strategy efficiently, enabling them to move away from laborious and unreliable counting strategies, such as 'counting all' and 'counting on'. Increasing fluency in efficient strategies will allow pupils to develop flexible and interlinked approaches to addition and subtraction. At a later stage, applying multiplication and division facts, rather than relying on skip-counting, will continue to develop flexibility with number.

Structures and contexts for calculations

There are multiple contexts (the word problem or real-life situation, within which a calculation is required) for each mathematical operation (i.e. addition) and, as well as becoming fluent with efficient calculating strategies, pupils also need to become fluent in identifying which operations are required. If they are not regularly exposed to a range of different contexts, pupils will find it difficult to apply their understanding of the four operations. For each operation, a range of contexts can be identified as belonging to one of the conceptual 'structures' defined below.

"In a technological age, in which most calculations are done on machines, it surely cannot be disputed that knowing which calculation to do is more important than being able to do the calculation."

Derek Haylock (2014); Mathematics Explained for Primary Teachers, p.56

The **structure** is distinct from both the **operation** required in a given problem and the **strategy** that may be used to solve the calculation. In order to develop good number sense and flexibility when calculating, children need to understand that many strategies (preferably the most efficient one for them!) can be used to solve a calculation, once the correct operation has been identified. There is often an implied link between the given structure of a problem context and a specific calculating strategy. Consider the following question: A chocolate bar company is giving out free samples of their chocolate on the street. They began the day with 256 bars and have given away 197. How many do they have remaining? The reduction context implicitly suggests the action of 'taking away' and might lead to a pupil, for example, counting back or using a formal algorithm to subtract 197 from 256 (seeing the question as $256 - 197 = \square$). However, it is much easier to find the difference between 197 and 256 by adding on (seeing the question as $197 + \square = 256$). Pupils with well-developed number sense and a clear understanding of the inverse relationship between addition and subtraction will be confident in manipulating numbers in this way.

Every effort is made to include multiple contexts for calculation in the Mathematics Mastery materials but, when teachers adapt the materials (which is absolutely encouraged), having an awareness of the different structures and being sure to include a range of appropriate contexts, will ensure that pupils continue to develop their understanding of each operation. The following list should not be considered to be exhaustive but defines the structures (and some suggested contexts) that are specifically included in the statutory objectives and the non-statutory guidance of the national curriculum. Specific structures and contexts are introduced in the Mathematics Mastery materials at the appropriate time, according to this guidance.

Importance of knowns vs unknowns and using part-whole understanding

One of the key strategies that pupils should use to identify the correct operation(s) to solve a given problem (in day-to-day life and in word problems) is to clarify the known and unknown quantities and identify the relationships between them. Owing to the inverse relationship between addition and subtraction, it is better to consider them together as 'additive reasoning', since changing which information is unknown can lead to either addition or subtraction being more suitable to calculate a solution for the same context. For the same reason, multiplication and division are referred to as 'multiplicative reasoning'. Traditionally, approaches involving key vocabulary have been the main strategy used to identify suitable operations but owing to the shared underlying structures, key words alone can be ambiguous and lead to misinterpretation (see for example the question below about Samir and Lena, where the key word 'less' might be identified, but addition is required to solve the problem).

A more effective strategy is to encourage pupils to establish what they know about the relationship between the known and unknown values and if they represent a part or the whole in the problem, supported through the use of part-whole models and/or bar models. In the structures exemplified

below, the knowns and unknowns have been highlighted. Where appropriate, the part-whole relationships have also been identified. Pupils should always be given opportunities to identify and discuss these, both when calculating and when problem-solving.

Standard and non-standard contexts

Using key vocabulary as a means of interpreting problems is only useful in what are in this document defined as 'standard' contexts, i.e. those where the language is aligned with the operation used to solve the problem. Take the following example:

*First there were 12 people on the bus. Then three **more** people got on. How many people are on the bus now?*

Pupils would typically identify the word 'more' and assume from this that they need to add the values together, which in this case would be the correct action. However, in non-standard contexts, identifying key vocabulary is unhelpful in identifying a suitable operation. Consider this question:

First there were 12 people on the bus and then some more people got on at the school. Now there are 15 people on the bus. How many people got on at the school?

Again the word 'more' would be identified, and a pupil may then erroneously add together 12 and 15. It is therefore much more helpful to consider known and unknown values and the relations between them.


Overexposure to standard contexts and lack of exposure to non-standard contexts will mean pupils are more likely to rely on 'key vocabulary' strategies, as they see that this works in most of the cases they encounter. It is therefore important, when adapting lesson materials, that non-standards contexts are used systematically, alongside standard contexts.

Additive reasoning

Change structures
augmentation (increasing)
 where an existing value has been added to

Standard

First there were 12 people on the bus. Then three more people got on. How many people are on the bus now?




"I know both parts. My first part is twelve and my second part is three. I don't know the whole. I need to add the parts of twelve and three to find the whole."

$$12 + 3 = ?$$

Non-standard

First there were 12 people on the bus and then some more people got on at the school. Now there are 15 people on the bus. How many people got on at the school?




"I know my first part is twelve and I know the whole is 15. I don't know the value of the second part. To find the second part, I could add on from 12 to make 15 or I could subtract 12 from 15."

$$12 + ? = 15 \quad 15 - 12 = ?$$

Non-standard

First there were some people on the bus then it stopped to pick up three more passengers at the bank. Altogether now there are 15 people on the bus. How many were people on the bus before it stopped at the bank?




"I know the value of the second part is three and that the whole is 15. I don't know the value of the first part. To find the first part, I could add on from three to make 15 or I could subtract three from 15."

$$? + 3 = 15 \quad 15 - 3 = ?$$

reduction (decreasing)
 where an existing value has been reduced

Standard

First Kieran had six plates in his cupboard. Then he took two plates out to use for dinner. How many plates are in the cupboard now?




"I know the whole is six. I know one of the part that has been taken away is two. I don't know the other part. I need to subtract the known part, two, from the whole, six, to find the remaining part."

$$6 - 2 = ? \quad 2 + ? = 6$$

Non-standard

First there were six plates in the cupboard. Then Kieran took some out for dinner. There are now four plates left in the cupboard. How many did Kieran take out?




"I know the whole is six and the remaining part is four. I don't know the part that was taken away. To find the part that was taken away I can add on from four to make six or I could subtract four from six."

$$6 - ? = 4 \quad 6 - 4 = ?$$

Non-standard

First there were some plates in the cupboard. Then Kieran took two out for dinner. Now there are four left. How many plates were in the cupboard to start with?



"I know the part that has been taken away is two and the part that is left is four. I don't know the whole. I can find the whole by adding the parts of four and two."

$$? - 2 = 4 \quad 2 + 4 = ?$$

Note: the 'first... then... now' structure is used heavily in KS1 to scaffold pupils' understanding of change structures. Once pupils are confident with the structures, such linguistic scaffolding can be removed, and question construction can be changed to expose pupils to a greater range of nuance in interpreting problems. For example, the second and third reduction problems could be reworded as follows:

Kieran took two plates out of his cupboard for dinner. There were four left. How many plates were in the cupboard to begin with?

There were six plates in the cupboard before Kieran took some out for dinner. If there were four plates left in the cupboard, how many did Kieran take out?

These present the same knowns and unknowns, and therefore the same bar models and resulting equations to solve the problems; however, the change in wording makes them more challenging to pupils who have only worked with a 'first... then... now' structure so far.

Part-whole structures


Combination (aggregation)/partitioning

combining two or more discrete quantities/splitting one quantity into two or more sub-quantities

Hakan and Sally have made a stack of their favourite books. Four books belong to Hakan, three to Sally. How many books are in the stack altogether?

"I know both parts. One part is four and the other part is three. I don't know the whole. I need to add the parts of three and four to find the whole."

$$4 + 3 = ? \quad 3 + 4 = ?$$




(Only one problem has been written for combination as, owing to the commutativity of addition, the only change in question wording would be to swap Hakan and Sally's names. The resulting bar model and calculation would be identical.)

Sally and Hakan have made a stack of their favourite books. There are seven books altogether. If three of them are Sally's, how many belong to Hakan?

"I know the whole is seven and that one of the parts is three. I don't know the other part. I need to add on from three to make seven or subtract three from seven to find the other part."


$$3 + ? = 7 \quad 7 - 3 = ?$$



Sally and Hakan have made a stack of their favourite books. There are seven books altogether. If four of them are Hakan's, how many belong to Sally?

"I know the whole is seven and that one of the parts is four. I don't know the other part. I need to add on from four to make seven or subtract four from seven to find the other part."

$$4 + ? = 7 \quad 7 - 4 = ?$$



Note: all part-whole contexts are considered to be 'standard', as the language of part-whole is unambiguous.

Comparison structures

Comparison structures involve a relationship between two quantities; their relationship is expressed as a difference. The structures vary by which of the values are known/unknown (the larger quantity, the smaller quantity and/or their difference). Part-whole language is not used here because the context contains not one single 'whole', but instead two separate quantities and it is the relationship between them being considered. Comparison bar models are therefore used to model these structures, which are known to be the most challenging for pupils to interpret.


Smaller quantity and larger quantity are known (comparative difference)

Standard

Navin has saved £19 from his pocket money. Sara has saved £31 from her pocket money. How much **more** has Sara saved than Navin? **or** How much **less** has Navin saved than Sara?

"I know one quantity is 19 and the other quantity is 31. I don't know the difference. To find the difference I could add on from 19 to make 31 or I could subtract 19 from 31."

$$19 + ? = 31 \quad 31 - 19 = ?$$




Smaller quantity and difference are known (comparative addition)

Standard

Ella has six marbles. Robin has three **more** than Ella. How many marbles does Robin have?

"I know the smaller quantity is six. I know the difference is three. I don't know the larger quantity. To find the larger quantity I need to add three to six."

$$6 + 3 = ?$$




Non-standard

Samir and Lena are baking shortbread but Lena's recipe uses 15g **less** butter than Samir's. If Lena needs to use 25g of butter, how much does Samir need?

"I know the smaller quantity is 25. I know the difference between the quantities is 15. I don't know the larger quantity. To find the larger quantity I need to add 15 to 25."

$$? - 15 = 25 \quad 25 + 15 = ?$$




Larger quantity and difference are known (comparative subtraction)

Non-standard

Ella has some marbles. Robin has three **more** than Ella and he has nine marbles in total. How many marbles does Ella have?

"I know the larger quantity is nine. I know the difference between the quantities is three. I don't know the smaller quantity. To find the smaller quantity I need to add on from three to make nine or subtract three from nine."

$$? + 3 = 9 \quad 9 - 3 = ?$$




Standard

Samir's shortbread recipe uses 40g of butter. Lena's recipe uses 15g **less** butter. How much butter does Lena need?

"I know one quantity is 40. I know the difference between the quantities is 15. I don't know the smaller quantity but I know it is 15 less than 40. To find the smaller quantity, I need to subtract 15 from 40."

$$40 - 15 = ? \quad ? + 15 = 40$$




Multiplicative reasoning

Repeated grouping structures

repeated addition

groups (sets) of equal value are combined or repeatedly added

There are four packs of pencils. Each contains five pencils. How many pencils are there?



"I know there are four equal parts and that each part has a value of five. I don't know the value of the whole. To find the whole, I need to multiply four and five."


$$5 + 5 + 5 + 5 = ?$$

$$5 \times 4 = ?$$

repeated subtraction (grouping)

groups (sets) of equal value are partitioned from the whole or repeatedly subtracted

There are 12 counters. If each child needs three counters to play the game, how many children can play?




"I know the whole is twelve and that the value of each equal part is three. To find the number of equal parts, I need to know how many threes are in twelve."

$$3 \times ? = 12 \quad 12 \div 3 = ?$$

sharing (into equal groups)

the whole is shared into a known number (must be a positive integer) of equal groups (sets)

Share twelve counters equally between three children. How many counters does each child get?



"I know the whole is twelve and the number of equal parts is three. I don't know the value of each part. To find the value of each part, I need to know what goes into twelve three times."


$$? \times 3 = 12 \quad 12 \div 3 = ?$$

Cartesian product of two measures

correspondence

calculating the number of unique combinations that can be created from two (or more) sets

Robin has three different hats and four different tops. How many different outfits can he create, that combine one hat and one top?



"I know how many hats there are and I know how many tops there are. I don't know the number of different outfits that can be created. To find the number of outfits, I need to find how many different tops can be worn with each hat or how many different hats can be worn with each top."

$$4 \times 3 = ? \quad 3 \times 4 = ?$$


Scaling structures

scaling up

('times greater/times as much')

the original value is increased by a given scale factor

Rita receives £2 pocket money every week. Sim earns ten times as much money for her paper round. How much money does Sim earn?



"I know one value is two and I know the second value is ten times greater. I don't know the second value. To find the second value, I need to multiply two by ten."


$$2 \times 10 = ?$$

scaling down

('times smaller/times less')

the original value is reduced by a given scale factor

The house in my model village needs to be half the height of the church. If the church is 8 cm tall, how tall does the house need to be?



"I know one value is eight and I know the second value is half as great. I don't know the second value. To find the second value, I need to halve eight (or divide it by two)."


$$\text{Half of 8 is ?}$$

$$8 \div 2 = ?$$

scaling up ('times as many')

the value of the original quantity is increased by a given scale factor

The Albert Hall can hold five times as many people as the Festival Hall. If the Festival Hall holds 1000 people, how many does the Albert Hall hold?




"I know one value is 1000 and I know the second value is five times greater. I don't know the second value. To find the second value, I need to multiply 1000 by five."

$$1000 \times 5 = ?$$

scaling down ('times fewer')

the value of the original quantity is decreased by a given scale factor

Anouska's garden pond has ten times fewer frogs than fish. If there are fifty fish, how many frogs are there?



"I know one value is 50 and I know the second value is ten times less. I don't know the second value. To find the second value, I need to divide fifty by ten."

$$50 \div 10 = ?$$

Progression in calculations

Year 1

National curriculum objectives linked to addition and subtraction

These objectives are explicitly covered through the strategies outlined in this document:







- Add and subtract one-digit and two-digit numbers to 100, including zero (N.B. Year 1 N.C. objective is to do this with numbers to 20).
- Add and subtract numbers using concrete objects, pictorial representations, and mentally, including: a two-digit number and ones, a two-digit number and tens, 2 two-digit numbers; add 3 one-digit numbers (Year 2).
- Represent and use number bonds and related subtraction facts within 20.
- Given a number, identify 1 more and 1 less.
- Show that addition of two numbers can be done in any order (commutative) but subtraction of one number from another cannot (Year 2).
- Recognise the inverse relationship between addition and subtraction and use this to solve missing number problems (Year 2).

The following objectives should be planned for lessons where new strategies are being introduced and developed:

- Read, write and interpret mathematical statements involving addition (+), subtraction (−) and equal (=) signs.
- Solve one-step problems that involve addition and subtraction, using concrete objects and pictorial representations, and missing number problems, such as $7 = \square - 9$.
- Solve problems with addition and subtraction:
 - Using concrete objects and pictorial representations, including those involving numbers, quantities and measures
 - Applying their increasing knowledge of mental methods


Teachers should refer to the definitions and guidance on the [structures for addition and subtraction](#) to provide a range of appropriate real-life contexts for calculations.

Y1 Addition

Strategy & guidance	CPA	
<p>Count all</p> <p><i>Joining two groups and then recounting all objects using one-to-one correspondence</i></p>	<p>$3 + 4 = 7$</p> 	 <p>$5 + 3 = 8$</p>
<p>Counting on</p> <p><i>As a strategy, this should be limited to adding small quantities only (1, 2 or 3) with pupils understanding that counting on from the greater number is more efficient.</i></p>	<p>$8 + 1 = 9$</p> 	<p>$15 = 12 + 3$</p> 
<p>Part-part-whole</p> <p><i>Teach both addition and subtraction alongside each other, as pupils will use this model to identify the inverse relationship between them.</i></p> <p><i>This model begins to develop the understanding of the commutativity of addition, as pupils become aware that the parts will make the whole in any order.</i></p>	  <p>$6 + 4 = 10$</p> <p>$10 = 6 + 4$</p> <p>$10 - 6 = 4$</p> <p>$10 - 4 = 6$</p> <p>$10 = 4 + 6$</p>	

**Regrouping ten
ones to make ten**

*This is an essential skill
that will support
column addition later
on.*



$$3 + 9 =$$

$$3 + 9 = 12$$




'Make ten' strategy

*Pupils should be
encouraged to start at
the greater number
and partition the
smaller number to
make ten.*



*The colours of the
beads on the bead
string make it clear
how many more need
to be added to make
ten.*

*Also, the empty spaces
on the ten frame make
it clear how many
more are needed to
make ten.*

$$6 + 5 = 11$$



$$4 + 9 = 13$$



Adding 1, 2, 3 more

Here the emphasis should be on the language rather than the strategy. As pupils are using the beadstring, ensure that they are explaining using language such as;

'1 more than 5 is equal to 6.'


'2 more than 5 is equal to 7.'

'8 is 3 more than 5.'

Over time, pupils should be encouraged to rely more on their number bonds knowledge than on counting strategies.



1

more than 5 $5 + 1 = 6$



2

more than 5 $5 + 2 = 7$



Adding three single digit numbers (make ten first)

Pupils may need to try different combinations before they find the two numbers that make 10.

The first bead string shows 4, 7 and 6. The colours of the bead string show that it makes more than ten.

The second bead string shows 4, 6 and then 7.

The final bead string shows how they have now been put together to find the total.




$$\begin{aligned} 4 + 7 + 6 &= 10 + 7 \\ &= 17 \end{aligned}$$

**Partitioning to add
(no regrouping)**

Place value grids and Dienes blocks could be used as shown in the diagram before moving onto pictorial representations. Dienes blocks should always be available, as the main focus in Year 1 is the concept of place value rather than mastering the procedure.

When not regrouping, partitioning is a mental strategy and does not need formal recording in columns. This representation prepares them for using column addition with formal recording.

$$24 + 13 = 37$$




$$24 + 13 = 37$$

**Introducing column
method for
addition,
regrouping only**

Dienes blocks and place value grids should be used as shown in the diagrams. Even when working pictorially, pupils should have access to Dienes blocks.

See additional guidance on unit pages for extra guidance on this strategy.

$$24 + 17$$



Tens Ones

$$\begin{array}{r} 24 \\ + 17 \\ \hline 41 \end{array}$$

First add the ones.
Re-group 10 ones to 1 ten.
Next add the tens.





Adding multiples of ten

Using the vocabulary of 1 ten, 2 tens, 3 tens etc. **alongside** 10, 20, 30 is important, as pupils need to understand that it is a **ten** and not a one that is being added and they need to understand that a '2' digit in the tens column has a value of twenty.


It also emphasises the link to known number facts. E.g. '2 + 3 is equal to 5. So 2 tens + 3 tens is equal to 5 tens.

$$50 = 30 + 20$$







$$3 \text{ tens} + 5 \text{ tens} = \underline{\quad} \text{ tens}$$

$$30 + 50 = \underline{\quad}$$



$$36 + 40 = \square$$

Y1 Subtraction

Strategy & guidance	CPA
<p>Taking away from the ones <i>When this is first introduced, the concrete representation should be based upon the diagram. Real objects should be placed on top of the images as one-to-one correspondence so that pupils can take them away, progressing to representing the group of ten with a tens rod and ones with ones cubes.</i></p>	 <p>$7 - 3 = 4$</p>  <p>$15 - 3 = 12$</p>  <p>$6 - 2 = 4$</p>  <p>$28 - 4 =$</p>
<p>Counting back <i>Subtracting 1, 2, or 3 by counting back</i></p> <p><i>Pupils should be encouraged to rely on number bonds knowledge as time goes on, rather than using counting back as their main strategy.</i></p>	 <p>$4 = 6 - 2$</p>  <p>$16 - 2 = 14$</p>



Part-part-whole

Teach both addition and subtraction alongside each other, as the pupils will use this model to identify the link between them. Pupils start with ten cubes placed on the whole.

They then remove what is being taken away from the whole and place it on one of the parts.

The remaining cubes are the other part and also the answer. These can be moved into the second part space.

$$10 - 6 = 4$$





Make ten strategy

To subtract a 1-digit number from a 2-digit number.


Pupils identify how many need to be taken away to make ten first, partitioning the number being subtracted. Then they take away the rest to reach the answer.

$$14 - 5 = 9$$



Regroup a ten into 10 ones

After the initial introduction, the Dienes blocks should be placed on a place value chart to support place value understanding. This will support pupils when they later use the column method.





$$20 - 4 =$$

Taking away from the tens

Pupils should identify that they can also take away from the tens and get the same answer.

This reinforces their knowledge of number bonds to 10 and develops their application of number bonds for mental strategies.

$$9 = 15 - 6$$



Partitioning to subtract without regrouping

Dienes blocks on a place value chart (developing into using images on the chart) could be used, as when adding 2-digit numbers, reinforcing the main concept of place value for Year 1.

When not regrouping, partitioning is a mental strategy and does not need formal recording in columns. This representation prepares them for using column subtraction with formal recording.

$$34 - 13 = 21$$





$$34 - 13 = 21$$

Subtracting multiples of ten

Using the vocabulary of 1 ten, 2 tens, 3 tens etc. alongside 10, 20, 30 is important as pupils need to understand that it is a **ten** not a one that is being taken away.

$$40 = 60 - 20$$



$$6 \text{ tens} - 2 \text{ tens} = \text{ } \text{tens}$$

$$60 - 20 = \text{ }$$

$$38 - 10 = 28$$



$$38 - 10 = \square$$




Column method with regrouping

This example shows how pupils should work practically when being introduced to this method.

There is no formal recording in columns in Year 1 but this practical work will prepare pupils for formal methods in Year 2.

See additional guidance on unit pages to support with this method.

$$34 - 17 = 17$$




National Curriculum objectives linked to multiplication and division

These objectives are explicitly covered through the strategies outlined in this document:

- Solve one-step problems involving multiplication and division, by calculating the answer using concrete objects, pictorial representations and arrays with the support of the teacher.

Teachers should refer to definitions and guidance on the [structures for multiplication and division](#) to provide a range of appropriate real-life contexts for calculations.



Y1 Multiplication

Strategy & guidance	CPA
<p>Skip counting in multiples of 2, 5, 10 from zero</p> <p><i>The representation for the amount of groups supports pupils' understanding of the written equation. So two groups of 2 are 2, 4. Or five groups of 2 are 2, 4, 6, 8, 10.</i></p> <p><i>Count the groups as pupils are skip counting.</i></p> <p><i>Number lines can be used in the same way as the bead string.</i></p> <p><i>Pupils can use their fingers as they are skip counting.</i></p>	 <p style="text-align: center;">$4 \times 5 = 20$</p>  <p style="text-align: center;">$2 \times 4 = 8$</p>
<p>Making equal groups and counting the total</p> <p><i>How this would be represented as an equation will vary. This could be 2×4 or 4×2. The importance should be placed on the vocabulary used alongside the equation. So this picture could represent 2 groups of 4 or 4 twice.</i></p>	  <p style="text-align: center;">Draw  to show $2 \times 3 = 6$</p>

Solve multiplications using repeated addition

This strategy helps pupils make a clear link between multiplication and division as well as exemplifying the 'repeated addition' structure for multiplication. It is a natural progression from the previous 'count all' strategy as pupils can be encouraged to 'count on'. However, as number bonds knowledge grows, pupils should rely more on these important facts to calculate efficiently.

$$3 \times 3 = 3 + 3 + 3$$



How many apples are there altogether?

$$3 + 3 + 3 = 9$$

Y1 Division

Strategy & guidance


Sharing objects into groups

Pupils should become familiar with division equations through working practically.


The division symbol is not formally taught at this stage.

CPA

$$10 \div 2 = 5$$



There are 10 sweets. Ring groups of 2.



There are _____ groups of 2.

Draw an equal number of apples for each basket.



There are five apples in each basket.

Progression in calculations

Year 2

National Curriculum objectives linked to addition and subtraction

These objectives are explicitly covered through the strategies outlined in this document:









- Add and subtract numbers using concrete objects, pictorial representations, and mentally, including: a two-digit number and ones; a two-digit number and tens; 2 two-digit numbers; adding three one-digit numbers.
- Add and subtract numbers mentally, including: a three-digit number and ones; a three-digit number and tens; a three-digit number and hundreds (Year 3).
- Recall and use addition and subtraction facts to 20 fluently, and derive and use related facts up to 100.
- Find 10 or 100 more or less than a given number (Year 3).
- Show that addition of two numbers can be done in any order (commutative) but subtraction of one number from another cannot.
- Recognise and use the inverse relationship between addition and subtraction and use this to check calculations and solve missing number problems.
- Add and subtract numbers with up to three digits, using formal written methods of columnar addition and subtraction (Year 3).







The following objectives should be planned for lessons where new strategies are being introduced and developed:

- Solve problems with addition and subtraction: using concrete objects and pictorial representations, including those involving numbers, quantities and measures; apply increasing knowledge of mental and written methods.
- Solve problems, including missing number problems, using number facts, place value and more complex addition and subtraction. (Year 3)



Teachers should refer to the definitions and guidance on the [structures for addition and subtraction](#) to provide a range of appropriate real-life contexts for calculations.


Y2 Addition



Strategy & guidance	CPA
<p>Part-part-whole</p> <p><i>Pupils explore the different ways of making 20. They can do this with all numbers using the same representations.</i></p> <p><i>This model develops knowledge of the inverse relationship between addition and subtraction and is used to find the answer to missing number problems.</i></p>	<div style="display: flex; justify-content: space-between; align-items: flex-start;"> <div style="text-align: center;">  </div> <div style="text-align: right;"> $20 = 17 + 3$ $20 = 3 + 17$ $20 - 3 = 17$ $20 - 17 = 3$ </div> </div> <div style="text-align: center; margin-top: 20px;">  </div> <div style="text-align: center; margin-top: 20px;">  </div> <div style="text-align: center; margin-top: 10px;"> $\square + \square = 20$ $20 - \square = \square$ $\square + \square = 20$ $20 - \square = \square$ </div> <hr style="width: 50%; margin: 20px auto;"/> <div style="text-align: center; margin-top: 10px;"> $\square + 1 = 16$ $16 - 1 = \square$ $1 + \square = 16$ $16 - \square = 1$ </div> <div style="text-align: center; margin-top: 20px;">  </div>
<p>Counting on in tens and hundreds</p>	<div style="display: flex; justify-content: space-around; align-items: center;">    </div> <div style="text-align: center; margin-top: 20px;">  </div> <div style="text-align: center; margin-top: 20px;">  </div>

Strategy & guidance	CPA												
<p>Make ten strategy</p>  <p><i>How pupils choose to apply this strategy is up to them; however, the focus should always be on efficiency.</i></p> <p><i>It relies on an understanding that numbers can be partitioned in different ways in order to easily make a multiple of ten.</i></p>	 $38 + 15 =$  												
<p>Partitioning to add without regrouping</p> <p><i>As in Year 1, this is a mental strategy rather than a formal written method. Pupils use the Dienes blocks (and later, images) to represent 3-digit numbers but do not record a formal written method if there is no regrouping.</i></p>	 $455 + 103 = 558$												
<p>Column method with regrouping</p> <p><i>Dienes blocks should be used alongside the pictorial representations; they can be placed on the place value grid before pupils make pictorial representations.</i></p> <p><i>As in Year 1, the focus for the column method is to develop a strong understanding of place value.</i></p>	<table border="1" data-bbox="598 1478 917 1702"> <thead> <tr> <th>hundreds</th> <th>tens</th> <th>ones</th> </tr> </thead> <tbody> <tr> <td>3</td> <td>5</td> <td>8</td> </tr> <tr> <td></td> <td>3</td> <td>7</td> </tr> <tr> <td>3</td> <td>9</td> <td>5</td> </tr> </tbody> </table> 	hundreds	tens	ones	3	5	8		3	7	3	9	5
hundreds	tens	ones											
3	5	8											
	3	7											
3	9	5											

Y2 Subtraction

Strategy & guidance	CPA
<p>Counting back in multiples of ten and one hundred</p>	 <p style="text-align: center;"> $75 \quad 85 \quad 95$ $-10 \quad -10$ $750 \quad 850 \quad 950$ $-100 \quad -100$ </p>
<p>Using known number facts to create derived facts</p> <p><i>Dienes blocks should be used alongside pictorial and abstract representations when introducing this strategy, encouraging pupils to apply their knowledge of number bonds to add multiples of ten and 100.</i></p>	 <p style="text-align: right;"> $8 - 4 = 4$ <i>leads to</i> $80 - 40 = 40$ <i>leads to</i> $800 - 400 = 400$ </p>
<p>Subtracting tens and ones</p> <p><i>Pupils must be taught to partition the second number for this strategy as partitioning both numbers can lead to errors if regrouping is required.</i></p>	<p style="text-align: center;">$53 - 12 = 41$</p>  <p style="text-align: center;"> $41 \quad 43 \quad 53$ $-2 \quad -10$ </p>

Strategy & guidance	CPA
<p>Round and adjust (sometimes known as a compensating strategy)</p> <p><i>Pupils must be taught to round the number that is being subtracted.</i></p> <p><i>Pupils will develop a sense of efficiency with this method, beginning to identify when this method is more efficient than subtracting tens and then ones.</i></p>	 <p>Number line for $53 - 17 = 36$:</p> <p>Start at 53. A green arc shows adding 3 to reach 56. A black arc shows subtracting 20 to reach 36.</p> <p>$53 - 17 = 36$</p>
<p>Make ten</p> <p><i>How pupils choose to apply this strategy is up to them. The focus should always be on efficiency.</i></p> <p><i>It relies on an understanding that numbers can be partitioned in different ways in order to subtract to a multiple of ten.</i></p> <p><i>Pupils should develop an understanding that the parts can be added in any order.</i></p>	 <p>Partitioning 15 into 2 and 13:</p> $42 - 15 =$ <p>Tree diagram for 15:</p> <pre> 15 / \ 2 13 / \ 10 3 </pre> <p>Number line 1: $42 - 15 = 27$ (steps: $42 \rightarrow 40$ (-2), $40 \rightarrow 30$ (-10), $30 \rightarrow 27$ (-3))</p> <p>Number line 2: $42 - 15 = 27$ (steps: $42 \rightarrow 32$ (-10), $32 \rightarrow 30$ (-2), $30 \rightarrow 27$ (-3))</p>

Strategy & guidance	CPA												
<p>Partitioning to subtract without regrouping</p> <p><i>As in Year 1, the focus is to develop a strong understanding of place value and pupils should always be using concrete manipulatives alongside the pictorial.</i></p> <p><i>Formal recording in columns is unnecessary for this mental strategy. It prepares them to subtract with 3-digits when regrouping is required.</i></p>	<div style="text-align: center;">  <p style="text-align: center;">$263 - 121 = 142$</p> </div>												
<p>Column method with regrouping</p> <p><i>The focus for the column method is to develop a strong understanding of place value and concrete manipulatives should be used alongside.</i></p> <p><i>Pupils are introduced to calculations that require two instances of regrouping (initially from tens to one and then from hundreds to tens). E.g. $232 - 157$ and are given plenty of practice using concrete manipulatives and images alongside their formal written methods, ensuring that important steps are not missed in the recording.</i></p> <p><i>Caution should be exercised when introducing calculations requiring 'regrouping to regroup' (e.g. $204 - 137$) ensuring ample teacher modelling using concrete manipulatives and images.</i></p>	<div style="text-align: center;"> <table style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th style="padding: 0 10px;">hundreds</th> <th style="padding: 0 10px;">tens</th> <th style="padding: 0 10px;">ones</th> </tr> </thead> <tbody> <tr> <td style="text-align: center; vertical-align: middle;">1^3</td> <td style="text-align: center; vertical-align: middle;">4</td> <td style="text-align: center; vertical-align: middle;">17</td> </tr> <tr> <td></td> <td style="text-align: center; vertical-align: middle;">-</td> <td style="text-align: center; vertical-align: middle;">18</td> </tr> <tr> <td colspan="3" style="text-align: center; border-top: 1px solid black; border-bottom: 1px solid black;"> 129 </td> </tr> </tbody> </table>  </div>	hundreds	tens	ones	1^3	4	17		-	18	129		
hundreds	tens	ones											
1^3	4	17											
	-	18											
129													

National Curriculum objectives linked to multiplication and division

These objectives are explicitly covered through the strategies outlined in this document:



- Recall and use multiplication and division facts for the 2, 5 and 10 multiplication tables, including recognising odd and even numbers.
- Recall and use multiplication and division facts for the 3 and 4 multiplication tables (Year 3).
- Show that multiplication of two numbers can be done in any order (commutative) but division of one number by another cannot.



The following objectives should be planned for lessons where new strategies are being introduced and developed:






- Calculate mathematical statements for multiplication and division within the multiplication tables and write them using the multiplication (\times), division (\div) and equal ($=$) signs.
- Solve problems involving multiplication and division, using materials, arrays, repeated addition, mental methods and multiplication and division facts, including problems in context.

Teachers should refer to definitions and guidance on the [structures for multiplication and division](#) to provide a range of appropriate real-life contexts for calculations.





Y2 Multiplication

Strategy & guidance	CPA																																																																																				
<p>Skip counting in multiples of 2, 3, 4, 5, 10 from zero</p> <p><i>Pupils can use their fingers as they are skip counting, to develop an understanding of 'groups of'.</i></p> <p><i>Dot arrays can be used to create a visual representation for the different multiplication facts. Bead strings, groups of cubes (or unifix / multilink towers) provide useful concrete representations.</i></p>	<table border="1" style="margin-left: auto; margin-right: auto; border-collapse: collapse; text-align: center;"> <thead> <tr> <th style="width: 20px;"></th> <th style="width: 20px;">1</th> <th style="width: 20px;">2</th> <th style="width: 20px;">3</th> <th style="width: 20px;">4</th> <th style="width: 20px;">5</th> </tr> </thead> <tbody> <tr><td style="width: 20px;">0</td><td></td><td></td><td></td><td></td><td></td></tr> <tr><td>1</td><td>●</td><td>●</td><td>●</td><td>●</td><td>●</td></tr> <tr><td>2</td><td>●</td><td>●</td><td>●</td><td>●</td><td>●</td></tr> <tr><td>3</td><td>●</td><td>●</td><td>●</td><td>●</td><td>●</td></tr> <tr><td>4</td><td>●</td><td>●</td><td>●</td><td>●</td><td>●</td></tr> <tr><td>5</td><td>●</td><td>●</td><td>●</td><td>●</td><td>●</td></tr> <tr><td>6</td><td>●</td><td>●</td><td>●</td><td>●</td><td>●</td></tr> <tr><td>7</td><td>●</td><td>●</td><td>●</td><td>●</td><td>●</td></tr> <tr><td>8</td><td>●</td><td>●</td><td>●</td><td>●</td><td>●</td></tr> <tr><td>9</td><td>●</td><td>●</td><td>●</td><td>●</td><td>●</td></tr> <tr><td>10</td><td>●</td><td>●</td><td>●</td><td>●</td><td>●</td></tr> <tr><td>11</td><td>●</td><td>●</td><td>●</td><td>●</td><td>●</td></tr> <tr><td>12</td><td>●</td><td>●</td><td>●</td><td>●</td><td>●</td></tr> </tbody> </table> 		1	2	3	4	5	0						1	●	●	●	●	●	2	●	●	●	●	●	3	●	●	●	●	●	4	●	●	●	●	●	5	●	●	●	●	●	6	●	●	●	●	●	7	●	●	●	●	●	8	●	●	●	●	●	9	●	●	●	●	●	10	●	●	●	●	●	11	●	●	●	●	●	12	●	●	●	●	●
	1	2	3	4	5																																																																																
0																																																																																					
1	●	●	●	●	●																																																																																
2	●	●	●	●	●																																																																																
3	●	●	●	●	●																																																																																
4	●	●	●	●	●																																																																																
5	●	●	●	●	●																																																																																
6	●	●	●	●	●																																																																																
7	●	●	●	●	●																																																																																
8	●	●	●	●	●																																																																																
9	●	●	●	●	●																																																																																
10	●	●	●	●	●																																																																																
11	●	●	●	●	●																																																																																
12	●	●	●	●	●																																																																																
<p>Multiplication as repeated addition</p> <p><i>Pupils apply skip counting to help find the totals of repeated additions.</i></p>	 $5 + 5 + 5 + 5 + 5 + 5 + 5 + 5 = \square$     $4 \times 3 = \square$																																																																																				

Strategy & guidance	CPA
<p>Arrays to represent multiplication equations</p> <p><i>Concrete manipulatives and images of familiar objects begin to be organised into arrays and, later, are shown alongside dot arrays. It is important to discuss with pupils how arrays can be useful.</i></p> <p><i>Pupils begin to understand multiplication in a more abstract fashion, applying their skip counting skills to identify the multiples of the 2x, 5x and 10x tables.</i></p> <p><i>The relationship between multiplication and division also begins to be demonstrated.</i></p>	
<p>Multiplication is commutative</p> <p><i>Pupils should understand that an array and, later, bar models can represent different equations and that, as multiplication is commutative, the order of the multiplication does not affect the answer.</i></p>	 <p>$3 \times 5 = \square$</p> <p>$5 \times 3 = \square$</p> <p>$12 = 3 \times 4$ $12 = 4 \times 3$</p>

Strategy & guidance	CPA										
<p>Use of part-part-whole model to establish the inverse relationship between multiplication and division</p> <p><i>This link should be made explicit from early on, using the language of the part-part-whole model, so that pupils develop an early understanding of the relationship between multiplication and division. Bar models (with Cuisenaire rods) should be used to identify the whole, the value of the parts and the number of parts.</i></p> <p><i>It is important to highlight that with multiplication, the parts are of equal value as this is different to how this model is used for addition and subtraction.</i></p>	<p>There are three equal parts. Each part has a value of three. What is the whole?</p> <div style="display: flex; justify-content: space-around;"> <div style="text-align: center;"> $3 \times 3 = \square$  </div> <div style="text-align: center;"> $9 \div 3 = \square$  </div> </div> <p>What multiplication and division equations can you write for each bar model? Prove that the equations are correct using a bead string.</p> <div style="display: flex; justify-content: center; align-items: center;"> <div style="text-align: center;">  </div> <div style="border: 1px solid black; padding: 5px; margin-left: 20px;"> <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="border: 1px solid black; width: 30px; height: 20px;"></td> <td style="text-align: center; padding: 0 5px;">×</td> <td style="border: 1px solid black; width: 30px; height: 20px;"></td> <td style="text-align: center; padding: 0 5px;">=</td> <td style="border: 1px solid black; width: 30px; height: 20px;"></td> </tr> <tr> <td style="border: 1px solid black; width: 30px; height: 20px;"></td> <td style="text-align: center; padding: 0 5px;">÷</td> <td style="border: 1px solid black; width: 30px; height: 20px;"></td> <td style="text-align: center; padding: 0 5px;">=</td> <td style="border: 1px solid black; width: 30px; height: 20px;"></td> </tr> </table> </div> </div>		×		=			÷		=	
	×		=								
	÷		=								
<p>Doubling to derive new multiplication facts</p> <p><i>Pupils learn that known facts from easier multiplication tables can be used to derive facts from related times tables using doubling as a strategy.</i></p> <p><i>At this stage they double the 2× table facts to derive the 4× table facts.</i></p>	<div style="display: flex; align-items: center; margin-bottom: 20px;">  <div style="margin-left: 20px;">$5 \times 2 = 10$</div> </div> <div style="display: flex; align-items: center;">  <div style="margin-left: 20px;">$5 \times 4 = 20$</div> </div>										

Y2 Division

Strategy & guidance	CPA
<p>Division as sharing</p> <p><i>Here, division is shown as sharing.</i></p> <p><i>If we have ten pairs of scissors and we share them between two pots, there will be 5 pairs of scissors in each pot.</i></p>	<p style="text-align: center;">$10 \div 2 = 5$</p>  
<p>Division as grouping</p> <p><i>Here, division is shown as grouping.</i></p> <p><i>If we have ten forks and we put them into groups of two, there are 5 groups.</i></p>	<p style="text-align: center;">$10 \div 2 = 5$</p>  

Strategy & guidance


Use of part-part-whole model to represent division equations and to emphasise the relationship between division and multiplication

Pupils use arrays of concrete manipulatives and images of familiar objects to solve division equations.

They begin to use dot arrays to develop a more abstract concept of division.


It is important to highlight that with multiplication and division, the parts are of equal value as this is different to how this model is used for addition and subtraction.

CPA



$$15 \div 5 = \boxed{3}$$

$$15 \div 3 = \boxed{5}$$



Write the division equations that the array represents.




$$20 \div 4 = \boxed{}$$

$$20 \div 5 = \boxed{}$$

The whole is nine. There are three equal parts. What is the value of each part?

$$9 \div 3 = \boxed{}$$



Progression in calculations

Year 3

National Curriculum objectives linked to addition and subtraction

These objectives are explicitly covered through the strategies outlined in this document:


- add and subtract numbers mentally, including:
 - a three-digit number and ones
 - a three-digit number and tens
 - a three-digit number and hundreds
- add and subtract numbers with up to four digits, using formal written methods of columnar addition and subtraction (four digits is Year 4)
- find 10 or 100 more or less than a given number
- find 1 000 more or less than a given number (Year 4)
- estimate the answer to a calculation and use inverse operations to check answers


The following objectives should be planned for lessons where new strategies are being introduced and developed:


- solve problems, including missing number problems, using number facts, place value, and more complex addition and subtraction

Teachers should refer to definitions and guidance on the [structures for addition and subtraction](#) to provide a range of appropriate real-life contexts for calculations.

Y3 Addition & Subtraction

Strategy & guidance	CPA														
<p>Add and subtract numbers mentally, including:</p> <ul style="list-style-type: none"> • a three-digit number and ones; • a three-digit number and tens; • a three-digit number and hundreds <p><i>Pupils learn that this is an appropriate strategy when they are able to use known and derived number facts or other mental strategies to complete mental calculations with accuracy.</i></p> <p><i>To begin with, some pupils will prefer to use this strategy only when there is no need to regroup, using number facts within 10 and derivations. More confident pupils might choose from a range of mental strategies that avoid written algorithms, including (but not exhaustively):</i></p> <ul style="list-style-type: none"> • known number facts within 20, • derived number facts, • ‘Make ten’, • round and adjust <p><i>See Year 2 guidance for exemplification of these – the use of concrete manipulatives other than Dienes blocks is important in reinforcing the use of these strategies.</i></p> <p><i>It is important that pupils are given plenty of (scaffolded) practice at choosing their own strategies to complete calculations efficiently and accurately. Explicit links need to be made between familiar number facts and the calculations that they can be useful for and pupils need to be encouraged to aim for efficiency.</i></p>	<p>It is important to model the mental strategy using concrete manipulatives in the first instance and pupils should be able to exemplify their own strategies using manipulatives if required, with numbers appropriate to the unit they are working on (3-digit numbers in Units 1 & 4; 4-digit numbers in Unit 13). However, pupils should be encouraged to use known facts to derive answers, rather than relying on counting manipulatives or images.</p> <p><u>No regrouping</u></p> <table style="width: 100%; border: none;"> <tr> <td style="text-align: left;">$345 + 30$</td> <td style="text-align: right;">$274 - 50$</td> </tr> <tr> <td style="text-align: left;">$1128 + 300$</td> <td style="text-align: right;">$1312 - 300$</td> </tr> <tr> <td style="text-align: left;">$326 + 342$</td> <td style="text-align: right;">$856 - 724$</td> </tr> </table> <div style="display: flex; align-items: center; margin-top: 10px;">  <div style="margin-left: 10px; color: red;"> <p>I know $4 + 3 = 7$, so 4 tens plus 3 tens is equal to 7 tens. $345 + 30 = 375$.</p> </div> </div> <p><u>With some regrouping</u></p> <table style="width: 100%; border: none;"> <tr> <td style="text-align: left;">$416 + 25$</td> <td style="text-align: right;">$232 - 5$</td> </tr> <tr> <td style="text-align: left;">$383 + 130$</td> <td style="text-align: right;">$455 - 216$</td> </tr> <tr> <td style="text-align: left;">$611 + 194$</td> <td style="text-align: right;">$130 - 40$</td> </tr> <tr> <td style="text-align: left;">$1482 + 900$</td> <td style="text-align: right;">$2382 - 500$</td> </tr> </table>	$345 + 30$	$274 - 50$	$1128 + 300$	$1312 - 300$	$326 + 342$	$856 - 724$	$416 + 25$	$232 - 5$	$383 + 130$	$455 - 216$	$611 + 194$	$130 - 40$	$1482 + 900$	$2382 - 500$
$345 + 30$	$274 - 50$														
$1128 + 300$	$1312 - 300$														
$326 + 342$	$856 - 724$														
$416 + 25$	$232 - 5$														
$383 + 130$	$455 - 216$														
$611 + 194$	$130 - 40$														
$1482 + 900$	$2382 - 500$														

Strategy & guidance	CPA						
<p>Written column method for calculations that require regrouping with up to 4-digits</p> <p><i>Dienes blocks should be used alongside the pictorial representations during direct teaching and can be used by pupils both for support and challenge. Place value counters can also be introduced at this stage.</i></p> <p><i>This work revises and reinforces ideas from Key Stage 1, including the focus on place value – see Year 2 exemplification.</i></p> <p><i>Direct teaching of the columnar method should require at least one element of regrouping, so that pupils are clear about when it is most useful to use it. Asking them ‘Can you think of a more efficient method?’ will challenge them to apply their number sense / number facts to use efficient mental methods where possible.</i></p> <p><i>As in Year 2, pupils should be given plenty of practice with calculations that require multiple separate instances of regrouping. In Year 3 they become more familiar with calculations that require ‘regrouping to regroup’. Understanding must be secured through the considered use of manipulatives and images, combined with careful use of language.</i></p> <p><i>Pupils should be challenged as to whether this is the most efficient method, considering whether mental methods (such as counting on, using known number facts, round and adjust etc.) may be likelier to produce an accurate solution.</i></p> <p><i>Pupils requiring support might develop their confidence in the written method using numbers that require no regrouping.</i></p> <p><i>See Unit materials for extra guidance on this strategy.</i></p>	<p>As for the mental strategies, pupils should be exposed to concrete manipulatives modelling the written calculations and should be able to represent their written work pictorially or with concrete manipulatives when required.</p> <p>Again, they should be encouraged to calculate with known and derived facts and should not rely on counting images or manipulatives.</p>  <p>5 + 6 = 11 so I will have 11 ones which I regroup for 1 ten and 1 one.</p> <p><u>Regrouping (including multiple separate instances)</u></p> <table data-bbox="869 1276 1332 1456"> <tr> <td>672 + 136</td> <td>734 – 82</td> </tr> <tr> <td>468 + 67</td> <td>831 - 76</td> </tr> <tr> <td>275 + 386</td> <td>435 – 188</td> </tr> </table> <p><u>‘Regrouping to regroup’</u></p> <p>204 – 137</p> <p>1035 - 851</p>	672 + 136	734 – 82	468 + 67	831 - 76	275 + 386	435 – 188
672 + 136	734 – 82						
468 + 67	831 - 76						
275 + 386	435 – 188						

Strategy & guidance	CPA
<p>Find 10, 100 more or less than a given number</p> <p><i>As pupils become familiar with numbers up to 1000, place value should be emphasised and comparisons drawn between adding tens, hundreds (and, in the last unit of the Summer term, thousands), including use of concrete manipulatives and appropriate images.</i></p> <p><i>After initial teaching, this should be incorporated into transition activities and practised regularly.</i></p>	<p>$142 + 100 = 242$</p>  <p>The CPA diagrams illustrate the addition of 100 to 142. The top diagram shows 142 (1 hundred, 4 tens, 2 units) and 100 (1 hundred) being added to reach 242 (2 hundreds, 4 tens, 2 units). The bottom diagram shows 142 (1 hundred, 4 tens, 2 units) and 100 (1 hundred) being added to reach 242 (2 hundreds, 4 tens, 2 units).</p>



National Curriculum objectives linked to multiplication and division




These objectives are explicitly covered through the strategies outlined in this document:


- count from 0 in multiples of 4, 8, 50 and 100
- recall and use multiplication and division facts for the 3, 4, 6, and 8 multiplication tables
- write and calculate mathematical statements for multiplication and division using the multiplication tables that they know, including for two-digit numbers times one-digit numbers, using mental methods
- solve problems, including missing number problems, involving multiplication and division, including positive integer scaling problems and correspondence problems in which n objects are connected to m objects

Teachers should refer to definitions and guidance on the [structures for multiplication and division](#) to provide a range of appropriate real-life contexts for calculations.

Y3 Multiplication

Strategy & guidance	CPA	
<p>Doubling to derive new multiplication facts</p> <p><i>Pupils continue to make use of the idea that facts from easier times tables can be used to derive facts from related times tables using doubling as a strategy.</i></p> <p><i>This builds on the doubling strategy from Year 2.</i></p>	<p>$3 \times 3 = 9$</p> 	<p>$3 \times 6 = \text{double } 9 = 18$</p> 

Strategy & guidance	CPA
<p>Skip counting in multiples of 2, 3, 4, 5, 6, 8 and 10</p> <p><i>Rehearsal of previously learnt tables as well as new content for Year 3 should be incorporated into transition activities and practised regularly.</i></p>	
<p>Use of part-part-whole model with arrays and bar models to establish commutativity and inverse relationship between multiplication and division</p> <p><i>In these contexts pupils are able to identify all the equations in a fact family.</i></p>	
<p>Ten times greater</p> <p><i>Pupils's work on this must be firmly based on concrete representations – the language of ten times greater must be well modelled and understood to prevent the numerical misconception of 'adding a zero'.</i></p>	

Strategy & guidance	CPA
<p>Multiplying by 10 and 100</p> <p><i>Building on the ten times greater work, pupils use appropriate Dienes blocks and place value counters to multiply 2, 3, 4, 5 and 10 by 10, 100 and 1000.</i></p>	<p>$5 \times 1 = 5$ </p> <p>$5 \times 10 = 50$ </p> <p>$3 \times 1 = 3$ </p> <p>$3 \times 100 = 300$ </p>
<p>Using known facts for multiplying by multiples of 10 and 100</p> <p><i>Pupils' growing understanding of place value, allows them to make use of known facts to derive multiplications using powers of 10.</i></p> <p><i>It is important to use tables with which they are already familiar (i.e. not 7 or 9 tables in Year 3)</i></p>	<p>$5 = 1 \times 5$ </p> <p>$50 = 10 \times 5$ </p> <p>$500 = 100 \times 5$ </p> <p>$5000 = 1000 \times 5$ </p> <p> $3 \times 2 = 6$ $30 \times 2 = 60$ $300 \times 2 = 600$ $3000 \times 2 = 6000$ </p> 


Strategy & guidance

Multiplication of 2-digit numbers with partitioning (no regrouping)

Children should always consider whether partitioning is the best strategy – if it is possible to use strategies such as doubling (some may use doubling twice for $\times 4$), they need to choose the most efficient strategy.

Children may wish to make jottings, including a full grid as exemplified here – but grid method is not a formal method and its only purpose is to record mental calculations. This supports the development of the necessary mental calculating skills but does not hinder the introduction of formal written methods in Year 4. Concrete manipulatives are essential to develop understanding.



CPA






Now add the total number of tens and ones

×	10	2	×	10	2
3			3	30	6

$3 \times 12 = 36$

Strategy & guidance	CPA																						
<p>Multiplication of 2-digit numbers with partitioning (regrouping)</p> <p><i>Using concrete manipulatives and later moving to using images that represent them, supports pupils' early understanding, leading towards formal written methods in Year 4.</i></p> <p><i>Once again, this is a mental strategy, which they may choose to support with informal jottings, including a full grid, as exemplified here.</i></p> <p><i>Pupils must be encouraged to make use of their known multiplication facts and their knowledge of place value to calculate, rather than counting manipulatives.</i></p>	<table border="1" data-bbox="560 293 839 479"> <tr> <td>×</td> <td>10</td> <td>4</td> </tr> <tr> <td>3</td> <td></td> <td></td> </tr> <tr> <td></td> <td>30</td> <td>12</td> </tr> </table> <table border="1" data-bbox="544 551 855 752"> <tr> <td>×</td> <td>40</td> <td>5</td> </tr> <tr> <td>3</td> <td></td> <td></td> </tr> </table>	×	10	4	3				30	12	×	40	5	3			<table border="1" data-bbox="895 293 1158 472"> <tr> <td>×</td> <td>10</td> <td>4</td> </tr> <tr> <td>3</td> <td>30</td> <td>12</td> </tr> </table>  <p>$14 \times 3 = 42$</p> 	×	10	4	3	30	12
×	10	4																					
3																							
	30	12																					
×	40	5																					
3																							
×	10	4																					
3	30	12																					

Y3 Division

Strategy & Guidance	CPA									
<p>Dividing multiples of 10, 100 and 1000 by 10, 100 and 1000 using scaling down</p> <p><i>Pupils use the strategy of 'scaling down', representing numbers with concrete manipulatives and making the value ten times smaller.</i></p>	<table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th style="width: 33%;">hundreds</th> <th style="width: 33%;">tens</th> <th style="width: 33%;">ones</th> </tr> </thead> <tbody> <tr> <td style="text-align: center;"></td> <td style="text-align: center;"></td> <td style="text-align: center;"></td> </tr> <tr> <td></td> <td style="text-align: center;">  3 </td> <td style="text-align: center;">  3 0  3 </td> </tr> </tbody> </table> <div style="margin-left: 200px; margin-top: 20px;"> <p>$3 \times 10 = 30$</p> <p>$30 \div 10 = 3$</p> </div>	hundreds	tens	ones					 3	 3 0  3
hundreds	tens	ones								
										
	 3	 3 0  3								
<p>Dividing multiples of 10, 100 and 1000 by 10, 100 and 1000 using grouping</p> <p><i>Pupils divide by 10, 100 and 1000 by making groups of the divisor.</i></p>	<p>$500 \div 100 = \square$</p> <p>My whole is 500 and the value of the equal parts is 100. How many parts are there?</p> <table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th style="width: 25%;">thousands</th> <th style="width: 25%;">hundreds</th> <th style="width: 25%;">tens</th> <th style="width: 25%;">ones</th> </tr> </thead> <tbody> <tr> <td></td> <td style="text-align: center;">  5 0 0 </td> <td style="text-align: center;">0</td> <td style="text-align: center;">0</td> </tr> </tbody> </table>	thousands	hundreds	tens	ones		 5 0 0	0	0	
thousands	hundreds	tens	ones							
	 5 0 0	0	0							

Progression in calculations

Year 4

National curriculum objectives linked to addition and subtraction

These objectives are explicitly covered through the strategies outlined in this document:


- add and subtract numbers with up to four digits, using the formal written methods of columnar addition and subtraction where appropriate
- find 1 000 more or less than a given number
- estimate and use inverse operations to check answers to a calculation






N.B. There is no explicit reference to mental calculation strategies in the programmes of study for Year 4 in the national curriculum. However, with an overall aim for fluency, appropriate mental strategies should always be considered before resorting to formal written procedures, with the emphasis on pupils making their own choices from an increasingly sophisticated range of strategies.



The following objectives should be planned for lessons where new strategies are being introduced and developed:

















- solve addition and subtraction two-step problems in contexts, deciding which operations and methods to use and why
- solve simple measure and money problems involving fractions and decimals to two decimal places

Y4 Addition & Subtraction

Strategies & Guidance	CPA
<p>Count forwards and backwards in steps of 10, 100 and 1000 for any number up to 10 000.</p> <p><i>Pupils should count on and back in steps of ten, one hundred and one thousand from different starting points. These should be practised regularly, ensuring that boundaries where more than one digit changes are included.</i></p> <p>Count forwards and backwards in tenths and hundredths</p>	  <p>Pay particular attention to boundaries where regrouping happens more than once and so more than one digit changes.</p> <p>E.g. $990 + 10$ or $19.9 + 0.1$</p>
<p>Using known facts and knowledge of place value to derive facts.</p> <p>Add and subtract multiples of 10, 100 and 1000 mentally</p> <p><i>Pupils extend this knowledge to mentally adding and subtracting multiples of 10, 100 and 1000. Counting in different multiples of 10, 100 and 1000 should be incorporated into transition activities and practised regularly.</i></p>	 <p>$2 + 4 = 6$</p> <p>$20 + 40 = 60$</p> <p>$200 + 400 = 600$</p> <p>$2000 + 4000 = 6000$</p>
<p>Adding and subtracting by partitioning one number and applying known facts.</p> <p><i>By Year 4 pupils are confident in their place value knowledge and are calculating mentally both with calculations that do not require regrouping and with those that do.</i></p>	<p>See Y3 guidance on mental addition & subtraction, remembering that use of concrete manipulatives and images in both teaching and reasoning activities will help to secure understanding and develop mastery.</p>

Strategies & Guidance	CPA
<p>Round and adjust</p> <p><i>Pupils should recognise that this strategy is useful when adding and subtracting near multiples of ten. They should apply their knowledge of rounding.</i></p> <p><i>It is very easy to be confused about how to adjust and so visual representations and logical reasoning are essential to success with this strategy.</i></p> <p><i>Build flexibility by completing the same calculation in a different order.</i></p>	<p>$3527 + 296 = 3827 - 4$</p>  <p>Completing the same calculation but adjusting first:</p> <p>$3527 + 296 = 3523 + 300$</p>  <p>$4523 - 3997 = 523 + 3$</p>  <p>Completing the same calculation but adjusting first:</p> <p>$4523 - 3997 = 4526 - 4000$</p> 
<p>Near doubles</p> <p><i>Pupils should be able to double numbers up to 100 and use this to derive doubles for multiples of ten. These facts can be adjusted to calculate near doubles.</i></p>	<p>$1600 + 1598 = \text{double } 1600 - 2$</p> 

Strategies & Guidance	CPA																
<p>Written column methods for addition</p> <p><i>Place value counters are a useful manipulative for representing the steps of the formal written method. These should be used alongside the written layout to ensure conceptual understanding and as a tool for explaining.</i></p> <p><i>This method and the language to use are best understood through the tutorial videos found here on the toolkit.</i></p>	 <table border="1" data-bbox="1152 1048 1396 1279"> <tr><td>5</td><td>2</td><td>7</td><td>3</td></tr> <tr><td>+</td><td>5</td><td>4</td><td>1</td></tr> <tr><td colspan="4"><hr/></td></tr> <tr><td>5</td><td>8</td><td>1</td><td>4</td></tr> </table>	5	2	7	3	+	5	4	1	<hr/>				5	8	1	4
5	2	7	3														
+	5	4	1														
<hr/>																	
5	8	1	4														
<p>Written column methods for subtraction</p> <p><i>Place value counters are a useful manipulative for representing the steps of the formal written method. These should be used alongside the written layout to ensure conceptual understanding and as a tool for explaining.</i></p> <p><i>This method and the language to use are best understood through the tutorial videos on the toolkit.</i></p>	 <table border="1" data-bbox="1184 1550 1391 1765"> <tr><td>4²</td><td>2¹⁵</td><td>2</td></tr> <tr><td colspan="3"><hr/></td></tr> <tr><td>1</td><td>0</td><td>8</td></tr> <tr><td colspan="3"><hr/></td></tr> <tr><td>1</td><td>0</td><td>8</td><td>1</td></tr> </table>	4 ²	2 ¹⁵	2	<hr/>			1	0	8	<hr/>			1	0	8	1
4 ²	2 ¹⁵	2															
<hr/>																	
1	0	8															
<hr/>																	
1	0	8	1														

Strategies & Guidance	CPA												
<p>Calculating with decimal numbers</p> <p><i>Assign different values to Dienes equipment. If a Dienes 100 block has the value of 1, then a tens rod has a value of 0.1 and a ones cube has a value of 0.01. These can then be used to build a conceptual understanding of the relationship between these.</i></p> <p><i>Place value counters are another useful manipulative for representing decimal numbers.</i></p> <p><i>All of the calculation strategies for integers (whole numbers) can be used to calculate with decimal numbers.</i></p>	<div style="text-align: center;">  </div> <p>$24.2 + 13.4 =$</p> <table border="1" style="width: 100%; text-align: center; border-collapse: collapse;"> <thead> <tr> <th style="width: 33%;">Tens</th> <th style="width: 33%;">Ones</th> <th style="width: 33%;">tenths</th> </tr> </thead> <tbody> <tr> <td>  </td> <td>  </td> <td>  </td> </tr> <tr> <td>  </td> <td>  </td> <td>  </td> </tr> <tr> <td> </td> <td> </td> <td> </td> </tr> </tbody> </table>	Tens	Ones	tenths									
Tens	Ones	tenths											
													
													

National Curriculum objectives linked to multiplication and division









These objectives are explicitly covered through the strategies outlined in this document:

- count from 0 in multiples of 6, 7, 9, 25 and 1000
- recall and use multiplication and division facts for multiplication tables up to 12×12
- write and calculate mathematical statements for multiplication and division using the multiplication tables that they know, including for two-digit numbers times one-digit numbers, using mental and progressing to formal written methods
- recognise and use factor pairs and commutativity in mental calculations
- use place value, known and derived facts to multiply and divide mentally, including: multiplying by 0 and 1; dividing by 1; multiplying together three numbers
- multiply two-digit and three-digit numbers by a one-digit number using formal written layout
- find the effect of dividing a one- or two-digit number by 10 and 100, identifying the value of the digits in the answer as ones, tenths and hundredths.

The following objectives should be planned for lessons where new strategies are being introduced and developed:

- solve problems involving multiplying and adding, including using the distributive law to multiply two digit numbers by one digit, integer scaling problems and harder correspondence problems such as n objects are connected to m objects.

Y4 Multiplication

Strategies & Guidance	CPA																				
<p>Multiplying by 10 and 100</p> <p><i>When you multiply by ten, each part is ten times greater. The ones become tens, the tens become hundreds, etc.</i></p> <p><i>When multiplying whole numbers, a zero holds a place so that each digit has a value that is ten times greater.</i></p> <p><i>Repeated multiplication by ten will build an understanding of multiplying by 100 and 1000</i></p>	<table border="1" style="width: 100%; text-align: center;"> <thead> <tr> <th style="width: 25%;">thousands</th> <th style="width: 25%;">hundreds</th> <th style="width: 25%;">tens</th> <th style="width: 25%;">ones</th> </tr> </thead> <tbody> <tr> <td></td> <td></td> <td></td> <td></td> </tr> <tr> <td></td> <td></td> <td></td> <td> 3</td> </tr> <tr> <td></td> <td> 3</td> <td></td> <td></td> </tr> <tr> <td>3</td> <td>0</td> <td>0</td> <td>0</td> </tr> </tbody> </table> <p style="text-align: right; margin-right: 20px;">$3 \times 10 = 30$</p> <p style="text-align: right; margin-right: 20px;">$3 \times 100 = 300$</p> <p style="text-align: right; margin-right: 20px;">$3 \times 1000 = 3000$</p>	thousands	hundreds	tens	ones								 3		 3			3	0	0	0
thousands	hundreds	tens	ones																		
																					
			 3																		
	 3																				
3	0	0	0																		
<p>Using known facts and place value for mental multiplication involving multiples of 10 and 100</p> <p><i>Pupils use their growing knowledge of multiplication facts, place value and derived facts to multiply mentally.</i></p> <p><i>Emphasis is placed on understanding the relationship (10 times or 100 times greater) between a known number fact and one to be derived, allowing far larger 'fact families' to be derived from a single known number fact.</i></p> <p><i>Knowledge of commutativity (that multiplication can be completed in any order) is used to find a range of related facts.</i></p>	<p style="text-align: center;"> factor factor product $3 \times 7 = 21$ </p> <div style="text-align: center;">  </div> <p style="text-align: center;"> factor factor product $7 \times 3 = 21$ </p> <div style="display: flex; justify-content: space-around;"> <div style="text-align: center;">  </div> <div style="text-align: center;">  </div> </div> <div style="display: flex; justify-content: space-around; margin-top: 20px;"> <div style="text-align: center;">  </div> <div style="text-align: center;">  </div> </div> <table style="width: 100%; margin-top: 20px;"> <tr> <td style="width: 50%; text-align: center;">$30 \times 7 = 210$</td> <td style="width: 50%; text-align: center;">$300 \times 7 = 2100$</td> </tr> <tr> <td style="text-align: center;">$70 \times 3 = 210$</td> <td style="text-align: center;">$700 \times 3 = 2100$</td> </tr> <tr> <td style="text-align: center;">$7 \times 30 = 210$</td> <td style="text-align: center;">$7 \times 300 = 2100$</td> </tr> <tr> <td style="text-align: center;">$3 \times 70 = 210$</td> <td style="text-align: center;">$3 \times 700 = 2100$</td> </tr> </table>	$30 \times 7 = 210$	$300 \times 7 = 2100$	$70 \times 3 = 210$	$700 \times 3 = 2100$	$7 \times 30 = 210$	$7 \times 300 = 2100$	$3 \times 70 = 210$	$3 \times 700 = 2100$												
$30 \times 7 = 210$	$300 \times 7 = 2100$																				
$70 \times 3 = 210$	$700 \times 3 = 2100$																				
$7 \times 30 = 210$	$7 \times 300 = 2100$																				
$3 \times 70 = 210$	$3 \times 700 = 2100$																				

Strategies & Guidance

Multiplying by partitioning one number and multiplying each part

Pupils build on mental multiplication strategies and develop an explicit understanding of distributive law, which allows them to explore new strategies to make more efficient calculations.





As well as partitioning into tens and ones (a familiar strategy), they begin to explore compensating strategies and factorisation to find the most efficient solution to a calculation.

Distributive law


$a \times (b + c) = a \times b + a \times c$

CPA


14×6



34×6



$30 \times 6 + 4 \times 6$



Mental multiplication of three 1-digit numbers, using the associative law

Pupils first learn that multiplication can be performed in any order, before applying this to choose the most efficient order to complete calculations, based on their increasingly sophisticated number facts and place value knowledge.

Four pots each containing two flowers which each have seven petals. How many petals in total?



$(4 \times 2) \times 7$ or $4 \times (2 \times 7)$

Strategies & Guidance

Short multiplication of 3-digit number by 1-digit number





To begin with pupils are presented with calculations that require no regrouping or only regrouping from the ones to the tens. Their conceptual understanding is supported by the use of place value counters, both during teacher demonstrations and during their own practice.

With practice pupils will be able to regroup in any column, including from the hundreds to the thousands, including being able to multiply numbers containing zero and regrouping through multiple columns in a single calculation.

This method and the language to use are best understood through the tutorial videos found [here](#) on the toolkit.

CPA





Exemplification of this process is best understood through viewing the video tutorial



To calculate 241×3 , represent the number 241. Multiply each part by 3, regrouping as needed.

$$\begin{array}{r}
 241 \\
 \times \quad 3 \\
 \hline
 723 \\
 \hline
 1
 \end{array}$$

Y4 Division

Strategies & Guidance	CPA																												
<p>Dividing by 10 and 100</p> <p><i>When you divide by ten, each part is ten times smaller. The hundreds become tens and the tens become ones. Each digit is in a place that gives it a value that is ten times smaller.</i></p> <p><i>When dividing multiples of ten, a place holder is no longer needed so that each digit has a value that is ten times smaller. E.g. $210 \div 10 = 21$</i></p>	<table border="1" style="width: 100%; text-align: center;"> <thead> <tr> <th>thousands</th> <th>hundreds</th> <th>tens</th> <th>ones</th> </tr> </thead> <tbody> <tr> <td></td> <td></td> <td></td> <td></td> </tr> <tr> <td></td> <td></td> <td></td> <td></td> </tr> <tr> <td></td> <td></td> <td>3</td> <td>0</td> </tr> <tr> <td></td> <td></td> <td></td> <td></td> </tr> <tr> <td></td> <td>3</td> <td>0</td> <td>0</td> </tr> <tr> <td>3</td> <td>0</td> <td>0</td> <td>0</td> </tr> </tbody> </table> <p style="text-align: right;"> $30 \div 10 = 3$ $300 \div 100 = 3$ $3000 \div 1000 = 3$ $300 \div 10 = 30$ $3000 \div 100 = 30$ $3000 \div 10 = 300$ </p>	thousands	hundreds	tens	ones											3	0						3	0	0	3	0	0	0
thousands	hundreds	tens	ones																										
																													
																													
		3	0																										
																													
	3	0	0																										
3	0	0	0																										
<p>Derived facts</p> <p><i>Pupils use their growing knowledge of multiplication facts, place value and derived facts to multiply mentally.</i></p> <p><i>Understanding of the inverse relationship between multiplication and division allows corresponding division facts to be derived.</i></p>	<p style="text-align: center;">$21 \div 3 = 7$</p> <div style="text-align: center;">  </div> <p style="text-align: center;">$21 \div 7 = 3$</p> <div style="display: flex; justify-content: space-around;"> <div style="text-align: center;">  </div> <div style="text-align: center;">  </div> </div> <div style="display: flex; justify-content: space-around; margin-top: 20px;"> <div style="text-align: center;">  </div> <div style="text-align: center;">  </div> </div> <table style="width: 100%; margin-top: 20px;"> <tr> <td style="text-align: center;">$210 \div 7 = 30$</td> <td style="text-align: center;">$2100 \div 7 = 300$</td> </tr> <tr> <td style="text-align: center;">$210 \div 3 = 70$</td> <td style="text-align: center;">$2100 \div 3 = 700$</td> </tr> <tr> <td style="text-align: center;">$210 \div 30 = 7$</td> <td style="text-align: center;">$2100 \div 300 = 7$</td> </tr> <tr> <td style="text-align: center;">$210 \div 70 = 3$</td> <td style="text-align: center;">$2100 \div 700 = 3$</td> </tr> </table>	$210 \div 7 = 30$	$2100 \div 7 = 300$	$210 \div 3 = 70$	$2100 \div 3 = 700$	$210 \div 30 = 7$	$2100 \div 300 = 7$	$210 \div 70 = 3$	$2100 \div 700 = 3$																				
$210 \div 7 = 30$	$2100 \div 7 = 300$																												
$210 \div 3 = 70$	$2100 \div 3 = 700$																												
$210 \div 30 = 7$	$2100 \div 300 = 7$																												
$210 \div 70 = 3$	$2100 \div 700 = 3$																												



Strategies & Guidance
Short division of 4-digit numbers by 1-digit numbers

Pupils start with dividing 4-digit numbers by 2, 3 and 4, where no regrouping is required. Place value counters are used simultaneously in a place value chart, to develop conceptual understanding.

They progress to calculations that require regrouping in the hundreds or tens columns.


Pupils build on their conceptual knowledge of division to become confident with dividing numbers where the tens digit is smaller than the divisor, extending this to any digit being smaller than the divisor.

Exemplification of this method and the language to use are best understood through viewing the tutorial videos found [here](#) on the toolkit.


CPA
Division as sharing

Division as grouping

Division of a one- or two-digit number by 10 and 100, identifying the value of the digits in the answer as ones, tenths and hundredths

When you divide by ten, each part is ten times smaller. The tens become ones and the ones become tenths. Each digit is in a place that gives it a value that is ten times smaller.

$$24 \div 10 = 2.4$$



$$24 \div 100 = 0.24$$



Progression in calculations

Year 5 + Year 6

Year 5 and Year 6 are together because the calculation strategies used are broadly similar, with Year 6 using larger and smaller numbers. Any differences for Year 6 are highlighted in red.

National Curriculum objectives linked to integer addition and subtraction






These objectives are explicitly covered through the strategies outlined in this document:

- add and subtract numbers mentally with increasingly large numbers
- add and subtract whole numbers with more than 4 digits, including using formal written methods (columnar addition and subtraction)
- use negative numbers in context, and calculate intervals across zero
- perform mental calculations, including with mixed operations and large numbers
- use estimation to check answers to calculations and determine, in the context of a problem, an appropriate degree of accuracy

The following objectives should be planned for lessons where new strategies are being introduced and developed:

- use rounding to check answers to calculations and determine, in the context of a problem, levels of accuracy
- solve addition and subtraction multi-step problems in contexts, deciding which operations and methods to use and why
- solve problems involving addition, subtraction, multiplication and division and a combination of these, including understanding the meaning of the equals sign.

Y5 and Y6 Addition & Subtraction

Strategies & Guidance	CPA																		
<p>Count forwards or backwards in steps of powers of 10 for any given number up to 1 000 000</p> <p><i>Skip counting forwards and backwards in steps of powers of 10 (i.e. 10, 100, 1000, 10 000 and 100 000) should be incorporated into transition activities and practised regularly.</i></p> <p><i>In Year 5 pupils work with numbers up to 1 000 000 as well as tenths, hundredths and thousandths.</i></p> <p><i>In Year 6 pupils work with numbers up to 10 000 000.</i></p>	<p>Support with place value counters on a place value chart, repeatedly adding the same counter and regrouping as needed.</p> <table border="1" style="width: 100%; text-align: center; border-collapse: collapse;"> <tr> <td>Hundred Thousands</td> <td>Ten Thousands</td> <td>Thousands</td> <td>Hundreds</td> <td>Tens</td> <td>Ones</td> <td>• tenths</td> <td>hundredths</td> <td>thousandths</td> </tr> <tr> <td> </td> </tr> </table> <p>Counting sticks and number lines:</p>  <p style="margin-left: 40px;">9700 9800 9900</p>  <p style="margin-left: 40px;">0 0.1 0.2</p> <p>Pay particular attention to boundaries where regrouping happens more than once and so more than one digit changes. e.g. $9900 + 100 = 10\,000$ or $99\,000 + 1000 = 100\,000$</p>	Hundred Thousands	Ten Thousands	Thousands	Hundreds	Tens	Ones	• tenths	hundredths	thousandths									
Hundred Thousands	Ten Thousands	Thousands	Hundreds	Tens	Ones	• tenths	hundredths	thousandths											
<p>Using known facts and understanding of place value to derive</p> <p><i>Using the following language makes the logic explicit: I know three ones plus four ones is equal to seven ones. Therefore, three ten thousands plus four ten thousands is equal to seven ten thousands.</i></p> <p><i>In Year 5 extend to multiples of 10 000 and 100 000 as well as tenths, hundredths and thousandths.</i></p> <p><i>In Year 6 extend to multiples of one million.</i></p> <p><i>These derived facts should be used to estimate and check answers to calculations.</i></p>	<div style="display: flex; justify-content: space-between;"> <div style="text-align: center;">  <p>$3 + 4 = 7$</p> </div> <div style="text-align: center;">  <p>$30\,000 + 40\,000 = 70\,000$</p> </div> <div style="text-align: center;">  <p>$300\,000 + 400\,000 = 700\,000$</p> </div> </div> <div style="margin-top: 20px;"> <p>$20\,000 + 40\,000 = 60\,000$</p> <p>$40\,000 + 20\,000 = 60\,000$</p> <p>$60\,000 - 40\,000 = 20\,000$</p> <p>$60\,000 - 20\,000 = 40\,000$</p> </div> <div style="margin-top: 20px;">  </div> <div style="margin-top: 20px;">  <div style="margin-left: 20px;"> <p>$0.6 = 0.2 + 0.4$</p> <p>$0.6 = 0.4 + 0.2$</p> <p>$0.2 = 0.6 - 0.4$</p> <p>$0.4 = 0.6 - 0.2$</p> </div> </div>																		

Strategies & Guidance
Subtraction by partitioning and applying known facts.


Pupils can use this strategy mentally or with jottings as needed.

Pupils should be aware of the range of choices available when deciding how to partition the number that is to be subtracted.


Pupils have experience with these strategies with smaller numbers from previous years and so the focus should be on developing flexibility and exploring efficiency.

CPA
Partitioning into place value amounts (canonical partitioning):


$$75\,221 - 14\,300 = 75\,221 - 10\,000 - 4\,000 - 300$$




Represent pictorially with a number line, starting on the right and having the arrows jump to the left:











Develop understanding that the parts can be subtracted in any order and the result will be the same:






Partitioning in different ways (non-canonical partitioning):



Extend the 'Make ten' strategy (see guidance in Y1 or Y2) to count back to a multiple of 10.



Strategies & Guidance	CPA
<p>Calculate difference by “counting back”</p> <p><i>It is interesting to note that finding the difference is reversible. For example, the difference between 5 and 2 is the same as the difference between 2 and 5. This is not the case for other subtraction concepts.</i></p>	<p>$75\,221 - 14\,300$</p> <p>Place the numbers either end of a numberline and work out the difference between them. Select efficient jumps.</p>  <p>Finding the difference is efficient when the numbers are close to each other:</p> <p>$9012 - 8976$</p> 
<p>Calculate difference by “counting on”</p> <p><i>Addition strategies can be used to find difference.</i></p>	<p>$75\,221 - 14\,300$</p>  <p>Finding the difference is efficient when the numbers are close to each other</p> <p>$9012 - 8976$</p> 

Strategies & Guidance	CPA
<p>Round and adjust</p> <p>Addition and subtraction using compensation</p> <p><i>Pupils should recognise that this strategy is useful when adding and subtracting near multiples of ten. They should apply their knowledge of rounding.</i></p> <p><i>It is very easy to be confused about how to adjust and so visual representations and logical reasoning are essential to success with this strategy.</i></p>	<p>Addition</p>  <p>$54\ 128 + 9987 = 54\ 128 + 10\ 000 - 13 = 64\ 128 - 13$</p> <p>Pupils should realise that they can adjust first:</p>  <p>$54\ 128 + 9987 = 54\ 128 - 13 + 10\ 000 = 54\ 115 + 10\ 000$</p> <p>Subtraction</p>  <p>$78\ 051 - 9992 = 78\ 051 - 10\ 000 + 8 = 68\ 051 + 8$</p> <p>Pupils should realise that they can adjust first:</p>  <p>$78\ 051 - 4960 = 78\ 051 + 40 - 5000 = 78\ 692 - 5000$</p>
<p>Near doubles</p> <p><i>Pupils should be able to double numbers up to 100 and use this to derive doubles for multiples of ten as well as decimal numbers. These facts can be adjusted to calculate near doubles.</i></p>	<p>$160 + 170 = \text{double } 150 + 10 + 20$</p> <p>$160 + 170 = \text{double } 160 + 10$ or $160 + 170 = \text{double } 170 - 10$</p> <p>$2.5 + 2.6 = \text{double } 2.5 + 0.1$</p>

Strategies & Guidance	CPA
<p>Partition both numbers and combine the parts</p> <p><i>Pupils should be secure with this method for numbers up to 10 000, using place value counters or Dienes to show conceptual understanding.</i></p> <p><i>If multiple regroupings are required, then pupils should consider using the column method.</i></p>	<p>$7230 + 5310 = 12\ 000 + 500 + 40$</p> <p style="text-align: center;">$200 + 300 = 500$</p>  <p style="text-align: center;">$7000 + 5000 = 12000$ $30 + 10 = 40$</p> <p>Pupils should be aware that the parts can be added in any order.</p>
<p>Written column methods for addition</p> <p><i>In Year 5, pupils are expected to be able to use formal written methods to add whole numbers with more than four digits as well as working with numbers with up to three decimal places.</i></p> <p><i>Pupils should think about whether this is the most efficient method, considering if mental methods would be more effective.</i></p> <p><i>Continue to use concrete manipulatives alongside the formal method.</i></p> <p><i>When adding decimal numbers with a different number of decimal places, in order to avoid calculation errors, pupils should be encouraged to insert zeros so that there is a digit in every row. This is not necessary for calculation and these zeros are not place holders as the value of the other digits is not changed by it being placed.</i></p> <p><i>Exemplification of this method and the language to use are best understood through viewing the tutorial videos found here on the toolkit.</i></p>	<p>For this method start with the digit of least value because if regrouping happens it will affect the digits of greater value.</p>  <p>Combine the counters in each column and regroup as needed:</p>  <p>Decimal numbers:</p> 

Strategies & Guidance	CPA
<p>Written column methods for subtraction</p> <p><i>In Year 5, pupils are expected to be able to use formal written methods to subtract whole numbers with more than four digits as well as working with numbers with up to three decimal places.</i></p> <p><i>Pupils should be given plenty of practice with calculations that require multiple separate instances of regrouping.</i></p> <p><i>In Year 3 and 4 they become more familiar with calculations that require ‘regrouping to regroup’. Understanding must be secured through the considered use of manipulatives and images, combined with careful use of language.</i></p> <p><i>Pupils should think about if this is the most efficient method, considering whether mental strategies (such as counting on, using known number facts, compensation etc.) may be likelier to produce an accurate solution.</i></p> <p><i>Exemplification of this method and the language to use are best understood through viewing the tutorial videos found here on the toolkit.</i></p>	<div style="display: flex; align-items: flex-start;"> <div style="margin-right: 20px;"> $\begin{array}{r} 41362 \\ - 32243 \\ \hline \end{array}$ $\begin{array}{r} 3\cancel{1}13\cancel{5}\cancel{1}2 \\ - 32243 \\ \hline 9119 \end{array}$ </div> <div>   </div> </div> <p>The term regrouping should be the language used. You can use the terms ‘exchange’ with subtraction but it needs careful consideration.</p> <p>You can regroup 62 as 50 and 12 (5 tens and 12 ones) instead of 60 and 2 (6 tens and 12 ones).</p> <p>Or you can ‘exchange’ one of the tens for 10 ones resulting in 5 tens and 12 ones.</p> <p>If you have exchanged, then the number has been regrouped.</p>

Progression in calculations

Year 5 + Year 6

National Curriculum objectives linked to multiplication and division

These objectives are explicitly covered through the strategies outlined in this document:








- multiply and divide whole numbers by 10, 100 and 1000
- multiply numbers up to 4 digits by a one- or two-digit number using a formal written method, including long multiplication for two-digit numbers
- multiply and divide numbers mentally drawing upon known facts
- divide numbers up to 4 digits by a one-digit number using the formal written method of short division and interpret remainders appropriately for the context
- multiply multi-digit numbers up to 4 digits by a two-digit whole number using the formal written method of long multiplication
- divide numbers up to 4 digits by a two-digit whole number using the formal written method of long division, and interpret remainders as whole number remainders, fractions, or by rounding, as appropriate for the context
- divide numbers up to 4 digits by a two-digit number using the formal written method of short division where appropriate, interpreting remainders according to the context
- multiply one-digit numbers with up to two decimal places by whole numbers
- use written division methods in cases where the answer has up to two decimal places

The following objectives should be planned for lessons where new strategies are being introduced and developed:

- solve problems involving multiplication and division including using their knowledge of factors and multiples, squares and cubes
- solve problems involving addition, subtraction, multiplication and division and a combination of these, including understanding the meaning of the equals sign
- use their knowledge of the order of operations to carry out calculations involving the four operations
- solve addition and subtraction multi-step problems in contexts, deciding which operations and methods to use and why
- solve problems involving addition, subtraction, multiplication and division
- solve problems involving the relative sizes of two quantities where missing values can be found by using integer multiplication and division facts.

Y5 and Y6 Multiplication



Strategies & Guidance	CPA																																	
<p>Multiply and divide whole numbers and those involving decimals by 10, 100 and 1000</p> <p><i>Avoid saying that you “add a zero” when multiplying by ten and instead use the language of place holder.</i></p> <p><i>Use place value counters and charts to visualise and then notice what happens to the digits.</i></p>	<p>When you multiply by ten, each part is ten times greater. The ones become tens, the tens become hundreds, etc.</p> <p>When multiplying whole numbers, a zero holds a place so that each digit has a value that is ten times greater.</p> <p>$102.14 \times 10 = 1021.4$</p> <table border="1" style="width: 100%; text-align: center; border-collapse: collapse;"> <thead> <tr> <th style="width: 12.5%;">Thousands</th> <th style="width: 12.5%;">Hundreds</th> <th style="width: 12.5%;">Tens</th> <th style="width: 12.5%;">Ones</th> <th style="width: 12.5%;">tenths</th> <th style="width: 12.5%;">hundredths</th> </tr> </thead> <tbody> <tr> <td></td> <td>100 100 100</td> <td></td> <td>1 1</td> <td>0.01</td> <td>0.1 0.1 0.1 0.1</td> </tr> <tr> <td>1000 1000 1000</td> <td></td> <td>10 10</td> <td>1</td> <td>0.01 0.01 0.01 0.01</td> <td></td> </tr> </tbody> </table> <p>When you divide by ten, each part is ten times smaller. The hundreds become tens and the tens become ones. Each digit is in a place that gives it a value that is ten times smaller.</p> <p>When dividing multiples of ten, a place holder is no longer needed so that each digit has a value that is ten times smaller.</p> <p>E.g. $210 \div 10 = 21$</p> <p>$210.3 \div 10 = 21.03$</p> <table border="1" style="width: 100%; text-align: center; border-collapse: collapse;"> <thead> <tr> <th style="width: 12.5%;">Hundreds</th> <th style="width: 12.5%;">Tens</th> <th style="width: 12.5%;">Ones</th> <th style="width: 12.5%;">tenths</th> <th style="width: 12.5%;">hundredths</th> </tr> </thead> <tbody> <tr> <td>100 100</td> <td>10</td> <td></td> <td>0.1 0.1 0.1 0.1</td> <td></td> </tr> <tr> <td></td> <td>10 10</td> <td>1</td> <td></td> <td>0.01 0.01 0.01 0.01</td> </tr> </tbody> </table>	Thousands	Hundreds	Tens	Ones	tenths	hundredths		100 100 100		1 1	0.01	0.1 0.1 0.1 0.1	1000 1000 1000		10 10	1	0.01 0.01 0.01 0.01		Hundreds	Tens	Ones	tenths	hundredths	100 100	10		0.1 0.1 0.1 0.1			10 10	1		0.01 0.01 0.01 0.01
Thousands	Hundreds	Tens	Ones	tenths	hundredths																													
	100 100 100		1 1	0.01	0.1 0.1 0.1 0.1																													
1000 1000 1000		10 10	1	0.01 0.01 0.01 0.01																														
Hundreds	Tens	Ones	tenths	hundredths																														
100 100	10		0.1 0.1 0.1 0.1																															
	10 10	1		0.01 0.01 0.01 0.01																														











Strategies & Guidance	CPA
<p>Using known facts and place value to derive multiplication facts</p> <p><i>Emphasis is placed on understanding the relationship (10 times or 100 times greater) between a known number fact and one to be derived, allowing far larger 'fact families' to be derived from a single known number fact.</i></p> <p><i>Knowledge of commutativity is further extended and applied to find a range of related facts.</i></p> <p><i>Pupils should work with decimals with up to two decimal places.</i></p> <p><i>These derived facts should be used to estimate and check answers to calculations.</i></p>	<div style="display: flex; justify-content: space-around;"> <div style="text-align: center;">  <p>$2 \times 3 = 6$ $3 \times 2 = 6$</p> </div> <div style="text-align: center;">  </div> </div> <p>$2 \times 30 = 60$ $30 \times 2 = 60$</p> <div style="display: flex; justify-content: space-around;"> <div style="text-align: center;">  </div> <div style="text-align: center;">  <p>$2 \times 3 \times 10 = 60$</p> </div> </div> <p>$3 \times 20 = 60$ $20 \times 3 = 60$</p> <div style="display: flex; justify-content: space-around;"> <div style="text-align: center;">  </div> <div style="text-align: center;">  </div> </div> <p>$2 \times 300 = 600$ $300 \times 2 = 600$</p> <p>$2 \times 30 \times 10 = 600$ $20 \times 3 \times 10 = 600$</p> <div style="display: flex; justify-content: space-around;"> <div style="text-align: center;">  </div> <div style="text-align: center;">  </div> </div> <p>$3 \times 200 = 600$ $200 \times 3 = 600$</p> <p>$20 \times 30 = 600$ $30 \times 20 = 600$</p>

These are the multiplication facts pupils should be able to derive from a known fact







2 100 000		700 000 x 3	70 000 x 30	7000 x 300	700 x 3000	70 x 30 000	7 x 300 000
210 000		70 000 x 3	7000 x 30	700 x 300	70 x 3000	7 x 30 000	
21 000		7000 x 3	700 x 30	70 x 300	7 x 3000		
2100		700 x 3	70 x 30	7 x 300			
210		70 x 3	7 x 30				
21	=	7 x 3					
2.1		0.7 x 3	7 x 0.3				
0.21		0.07 x 3	0.7 x 0.3	7 x 0.03			
0.021		0.007 x 3	0.07 x 0.3	0.7 x 0.03	7 x 0.003		

Strategies & Guidance	CPA
<p>Doubling and halving</p> <p><i>Pupils should experience doubling and halving larger and smaller numbers as they expand their understanding of the number system.</i></p> <p><i>Doubling and halving can then be used in larger calculations.</i></p>	<div data-bbox="730 297 1262 504" style="text-align: center;"> </div> <p>Multiply by 4 by doubling and doubling again e.g. $16 \times 4 = 32 \times 2 = 64$</p> <p>Divide by 4 by halving and halving again e.g. $104 \div 4 = 52 \div 2 = 26$</p> <div data-bbox="643 790 1342 1025" style="text-align: center;"> </div> <p>Multiply by 8 by doubling three times e.g. $12 \times 8 = 24 \times 4 = 48 \times 2 = 96$</p> <p>Divide by 8 by halving three times e.g. $104 \div 8 = 52 \div 4 = 26 \div 2 = 13$</p> <div data-bbox="592 1319 1401 1487" style="text-align: center;"> </div> <p>Multiply by 5 by multiplying by 10 then halving, e.g. $18 \times 5 = 180 \div 2 = 90$.</p> <p>Divide by 5 by dividing by 10 and doubling, e.g. $460 \div 5 = \text{double } 46 = 92$</p>

Strategies & Guidance	CPA
<p>Multiply by partitioning one number and multiplying each part</p> <p>Distributive law</p> <p>$a \times (b + c) = a \times b + a \times c$</p> <p>Build on pupils' understanding of arrays of counters to represent multiplication to see that area models can be a useful representation:</p>	<p>$8 \times 14 = 8 \times 10 + 8 \times 4$</p>  <p>80 32</p> <p>Cuisenaire rods to build arrays</p>  <p>14</p> <p>8 80 32</p> <p>Represent with area model</p>  <p>10 x 8 4 x 8</p> <p>0 80 100 112</p> <p>Jottings on a number line</p> <p>Bead string where each bead has a value of 8:</p>  <p>80 32</p>
<p>Using knowledge of factors</p> <p><i>In Year 5 pupils are expected to be able to identify factor pairs and this knowledge can be used to calculate.</i></p> <p><i>Pupils will be using the commutative and associative laws of multiplication.</i></p> <p>Commutative law</p> <p>$a \times b = b \times a$</p> <p>Associative law</p> <p>$a \times b \times c = (a \times b) \times c$</p> <p>$= a \times (b \times c)$</p> <p><i>They should explore and compare the different options and choose the most efficient order to complete calculations.</i></p>	<p>Calculate 6×24 by using factor pairs of 24</p> <p>Two and twelve are factors of 24:</p> <p>$6 \times 2 \times 12$</p>  <p>2 12</p> <p>6 6</p> <p>Three and eight are factors of 24:</p> <p>$6 \times 3 \times 8$</p>  <p>3 8</p> <p>6 6</p> <p>Four and six are factors of 24:</p> <p>$6 \times 4 \times 6$</p>  <p>4 6</p> <p>6 6</p>

Strategies & Guidance	CPA																	
<p>Formal written method of short multiplication</p> <p><i>Conceptual understanding is supported by the use of place value counters, both during teacher demonstrations and during their own practice.</i></p> <p><i>Exemplification of this method and the language to use are best understood through viewing the tutorial videos found here on the toolkit.</i></p>	$\begin{array}{r} 241 \\ \times 3 \\ \hline 723 \\ \hline 1 \end{array}$	<table border="1" style="width: 100%; text-align: center;"> <thead> <tr> <th>Hundreds</th> <th>Tens</th> <th>Ones</th> </tr> </thead> <tbody> <tr> <td></td> <td></td> <td></td> </tr> <tr> <td>  </td> <td></td> <td></td> </tr> </tbody> </table>	Hundreds	Tens	Ones													
Hundreds	Tens	Ones																
																		
<p>Multiplying by a 2-digit number</p> <p>Formal written method of long multiplication</p> <p><i>In Year 6 pupils are extended from multiplication by a 1-digit number to multiplication by a 2-digit number.</i></p> <p><i>Extend the place value chart model used in Year 4, using an additional row on the place value chart.</i></p> <p><i>Extend understanding of the distributive law to develop conceptual understanding of the two rows of the formal written method.</i></p> <p><i>Dienes blocks can be used to construct area models to represent this.</i></p>	$\begin{array}{r} 243 \\ \times 12 \\ \hline 486 \\ 2430 \\ \hline \end{array}$	<table border="1" style="width: 100%; text-align: center;"> <thead> <tr> <th>Thousands</th> <th>Hundreds</th> <th>Tens</th> <th>Ones</th> </tr> </thead> <tbody> <tr> <td></td> <td></td> <td></td> <td></td> </tr> <tr> <td></td> <td>  </td> <td></td> <td></td> </tr> <tr> <td>  </td> <td></td> <td></td> <td></td> </tr> </tbody> </table> <p style="text-align: right;">243 x 2</p> <p style="text-align: right;">243 x 10</p> $\begin{array}{r} 34 \\ \times 12 \\ \hline 68 \\ 340 \\ \hline 408 \end{array}$ <p>12 x 34</p>  <p style="text-align: right;">10 x 34 = 340</p> <p style="text-align: right;">2 x 34 = 68</p>	Thousands	Hundreds	Tens	Ones												
Thousands	Hundreds	Tens	Ones															
																		
																		

Y5 and Y6 Division

Strategies & Guidance	CPA
<p>Deriving facts from known facts</p> <p><i>Pupils use their growing knowledge of multiplication facts, place value and derived facts to multiply mentally.</i></p> <p><i>Understanding of the inverse relationship between multiplication and division allows corresponding division facts to be derived.</i></p>	<p style="text-align: center;">CPA</p> <p> $6 \div 2 = 3$   $6 \div 3 = 2$ </p> <p> $60 \div 2 = 30$   $60 \div 3 = 20$ $60 \div 30 = 2$ </p> <p> $600 \div 2 = 300$   $600 \div 3 = 200$ $600 \div 300 = 2$ </p>
<p>Using knowledge of multiples to divide</p> <p><i>Using an area model to partition the whole into multiples of the divisor (the number you are dividing by).</i></p>	<p>  $112 \div 8 = 80 \div 8 + 32 \div 8$ </p> <p>   </p> <p> $1260 \div 6 = 1200 \div 6 + 60 \div 6$ </p> <p>   </p>

Strategies & Guidance	CPA
<p>Using knowledge of factors to divide</p> <p><i>Pupils explore this strategy when using repeated halving.</i></p> <p><i>$2 \times 2 = 4$ and so if you divide by 4 the same result can be achieved by dividing by two and then by two again.</i></p>	<div style="text-align: center;"> <p>24</p>  <p>144</p> <p>$144 \div 24$</p> </div> <div style="text-align: center; margin-top: 20px;"> <p>12 12</p>  <p>72 144</p> <p>$144 \div 2 \div 12$</p> </div> <p>I know 2 and 12 are a factor pair of 24 and so I can divide by 2 and then by 12.</p>

Strategies & Guidance
Short division
Dividing a 4-digit numbers by 1-digit numbers

The thought process of the traditional algorithm is as follows:

How many 4s in 8? 2

How many 4s in 5? 1 with 1 remaining so regroup.

How many 4s in 12? 3

How many 4s in 8? 2

Warning: If you simply apply place value knowledge to each step, the thinking goes wrong if you have to regroup.

How many 4s in 8000? 2000

How many 4s in 500?

100 with 1 remaining (illogical)

The answer would be 125.

Sharing the dividend builds conceptual understanding however doesn't scaffold the "thinking" of the algorithm.

Using place value counters and finding groups of the divisor for each power of ten will build conceptual understanding of the short division algorithm.


Area models are also useful representations, as seen with other strategies and exemplified for long division.

Exemplification of this method and the language to use are best understood through viewing the tutorial videos found [here](#) on the toolkit.

CPA

$8528 \div 4$

$$\begin{array}{r} 2132 \\ 4 \overline{) 8528} \end{array}$$

Sharing



8 thousands shared into 4 equal groups

5 hundreds shared into 4 equal groups

Regroup 1 hundred for 10 tens

12 tens shared into 4 equal groups

8 ones shared into 4 equal groups.

Grouping



How many groups of 4 thousands in 8 thousands?

How many groups of 4 hundreds in 5 hundreds?

Regroup 1 hundred for 10 tens.

How many groups of 4 tens in 12 tens?

How many groups of 4 ones in 8 ones?

Strategies & Guidance	CPA
<p>Long division</p> <p>Dividing a 4-digit number by a 2-digit number</p> <p><i>Follow the language structures of the short division strategy. Instead of recording the regrouped amounts as small digits the numbers are written out below. This can be easier to work with when dividing by larger numbers.</i></p> <p><i>If dividing by a number outside of their known facts, pupils should start by recording some multiples of that number to scaffold.</i></p>	<div style="text-align: right; margin-bottom: 20px;"> $\begin{array}{r} 34 \\ 12 \overline{) 408} \\ \underline{36} \\ 48 \\ \underline{48} \\ 0 \end{array}$ </div> <p>$408 \div 12$</p>  <p style="text-align: center;"> $30 \times 12 = 360$ $4 \times 12 = 48$ </p>



Mathematics Mastery vocabulary list

This document highlights the vocabulary introduced throughout the primary curriculum – from Reception to Year 6.

The vocabulary listed here is vocabulary that **pupils** are expected to use and understand on a daily basis within that year group, though the definitions are written for teacher reference and would not necessarily be shared with children as they stand. The vocabulary listed is cumulative and builds on the vocabulary previously introduced. Teachers should also consult with the Mathematics Mastery Primary Glossary.


This is a working document and will be updated as required.


Reception	Definition	Example
Above	Used to describe a higher position than another object.	The Maths Meetings board is above the sink.
Add	Carry out the process of addition.	I can add two numbers together to find a total.
Addition	The operation to combine at least two numbers or quantities to form a further number or quantity, the sum or total. Addition is the inverse operation to subtraction.	Three plus seven is equal to ten. This is an addition equation.
Altogether	In total.	That will be £2 altogether please.
Balance	A measuring tool used to weigh objects. It has two dishes hanging on a bar. Both dishes will be level when the contents weigh the same. Also, as a verb, indicates equivalence and equality.	The objects in the balance are unequal in weight because the dish on the right side is lower down than the dish on the left side. The two objects balance which means they have the same mass.
Before	In front of or prior to.	The number '3' comes before '5' on the number line.
Below	Used to describe a lower position than another object.	The sink is below the Maths Meetings board.
Between	Indicates a position in relation to two other places or objects on either side.	The teacher is standing between two tables.
Capacity	The amount of liquid a container can hold.	This cup is full to capacity because it cannot hold any more water.
Circle	The name of a 2-D shape. A circle has a curved side.	
Clock	A tool used to measure time.	The clock shows us that the time is now 2 o'clock.
Compare	Look for similarities and/or differences between at least two objects or sets.	I can compare these two sets – this set has more.
Corner	A point where two or more lines meet. The correct mathematical term is vertex (vertices).	The table has four corners (vertices).
Cost	A monetary value assigned to a good or service.	This apple costs 10p. What coin could I use to pay for it?


Count	Assigning one number name to each of a set of objects to determine how many there are.	I counted the children in the group – there are four so we will need four pencils.
Cube	A 3-D shape with six identical square faces.	
Cuboid	A 3-D shape with six rectangular faces.	
Curved surface	A non-plane surface of a 3-D shape. Both cones and cylinders have curved surfaces.	The cone has a curved surface .
Cylinder	A 3-D shape with two circular faces joined by a curved surface.	
2-D	Abbreviation for two-dimensional. A figure is two-dimensional if it lies on a plane.	A square is a 2-D shape.
3-D	Abbreviation for three-dimensional. A solid is three-dimensional and occupies space.	A cylinder is a 3-D shape.
Describe	To express mathematical features, qualities and details in words.	Can you describe the properties of a cube?
Difference	The numerical difference between two numbers or sets of objects. It is found by comparing the quantity of one set of objects with another.	The difference between ten and six is four.
Direction	The orientation of a line in space.	Which direction should we jump – forwards or backwards?
Distance	A measure between two points or things.	The distance between my house and the school is longer than that between the school and the train station.
Double	To multiply by two or add a value to itself.	Ten is double five.
Edge	A line segment joining two vertices of a plane figure (2-D shape) and the intersection of two plane faces (in a 3-D shape).	A triangle has three edges and a cube has 12 edges .
Empty	Containing nothing. Most commonly used in the context of measures.	There is no more water left in the jug – it is empty .
Equal	Indicates equivalence between two values and can be expressed with the symbol '='. The symbol is read as 'is equal to' which means the same as. Expressions on either side of the symbol have the same value.	My sets are equal because there are four bears in this set and there are four bears in this set.
Face	One of the plane surfaces of a solid shape.	A cube has six faces .



Fewer	A lesser amount – used when counting discrete objects, i.e. countable objects such as, pens, teddies, counters, etc.	There are fewer buttons on my coat than yours.
First	Comes before all others in time or position.	First I brush my teeth. Then I go to bed.
Flat	A level surface.	The table has a flat rectangular surface.
Full	Contains/holds as much or as many as possible; has no empty space.	The juice carton is not full because I drank some.
Group	To make equal size groups. This is one model for division.	I will group the crayons equally so that each person gets two.
Half	One of two equal parts of a shape, quantity or object.	I have shared the dolls into two equal groups – I have half and you have half .
Intersection of sets	Where the two subsets overlap in a Venn diagram. Objects or values which belong to both subsets are placed here.	The number 4 belongs in the intersection because it is even <i>and</i> less than 5.
Last	Comes after all others in time or order.	Rory is the last person in the line.
Length	A linear measurement.	The length of my snake is shorter than yours.
Less	A smaller amount or not as much.	I have 15p and you have 7p. you have less money than me.
Line	A set of adjacent points that has length but no width.	I have drawn a line matching the number four with the four ducks.
Long	An adjective used to describe length.	I have a long piece of string.
Mass	A measure relating to the amount of matter within a given object.	The mass of the school bag is greater than the mass of the book.
Measure	To find the size of something in a given unit.	How might we measure how much flour we need to bake a cake?
Minus	A name for the symbol '-', which denotes the operation of subtraction.	Three minus one is equal to two.
More	A greater amount.	I have six apples and you have two. I have more .
Next	Comes immediately after the present one in order.	The next shape in my pattern is a square.
Number bond	A pair of numbers with a given total.	Five and four make a number bond to nine.
Number line	A linear, continuous representation of number. Each number occupies a point on the line, and there is an equal interval between each number.	This number line starts at zero and ends at ten.
Number track	A linear, discrete representation of number. Each number is positioned in a square on the track.	I can count from one to ten, moving a counter along this number track .
Order	Describes the placement of items according to given criteria or in a pattern. As a verb, to place items according to given criteria or in a pattern.	I have ordered the bears from smallest to biggest.



Venn diagram	Two or more circles which represent given sets and intersect according to these.	
Vertex (pl. vertices)	The point at which two or more lines intersect.	This shape has five vertices.
Weight	The force exerted on an object by gravity. Weight therefore changes with a change in gravitational force. Used interchangeably with mass until KS2.	The weight of this book is heavier than the pencil.
Zero	The number before one. It is neither positive nor negative.	Zero comes before one on the number track.

Year 1	Definition	Example
Analogue clock	A clock with a face and hands.	
Anticlockwise	Movement in the opposite direction to the motion of the hands of a clock.	
Approximate	The number is not exact but it is close.	Our PSHE lesson lasts approximately half an hour.
Array	An arrangement of counters or numbers, in columns and rows, used to represent multiplication and division	 This array shows 3×4 , 4×3 , $12 \div 4$ and $12 \div 3$
Block graph	The pre-cursor to the bar graph, this representation of data has an x- and y-axis and one block represents one item. Each block is adjoined to the adjacent block.	
Chart	A table or graph.	I will mark one day for the sun on our weather chart .
Chronological	In time order.	I ordered the events in my day chronologically . I woke up, ate my breakfast, went to school then came home.
Clockwise	Movement in the direction of the hands of a clock.	

Cone	A 3-D shape with one circular plane face, which tapers to an apex.	
Continuous surface	An outer boundary of a 3-D object which is uninterrupted by any plane surfaces.	A sphere has a continuous surface .
Data	Quantitative information which has been counted or measured.	This block graph shows us data for the colour of the cars in the car park.
Decreasing	Becoming smaller in value. Used in relation to number sequences.	15, 14, 13, 12. This number pattern is decreasing by one each time.
Diagram	An illustration, drawing or representation.	I will draw a diagram to show how I programed my floor toy to move.
Digit	One of the ten Arabic numerals 0 to 9, from which we compose numbers.	The number 54 has the digit five in the tens column and the digit four in the ones. The digit five has a value of fifty.
Divide	To share or group into equal parts.	I can divide 12 by three using grouping or sharing.
Estimate	An appropriately accurate guess, depending on the context and numbers involved.	I estimate there are eight cubes in the cup because it looks about double four but fewer than ten.
Even number	A number with a 0, 2, 4, 6 or 8 in the ones and therefore exactly divisible by two.	32 is an even number .
Facts	Related to the four operations (+, -, ×, ÷). Pupils should be supported in achieving fluency, i.e. very fast recall, in these facts. These then become known facts .	Number bonds to and within 10 and 20 are facts, e.g. $3 + 7 = 10$.
Fraction	<ol style="list-style-type: none"> 1. A part of a whole number, quantity or shape. 2. Expressing a division relationship between two integers in the form $\frac{a}{b}$. 	I have shared my sweets into four equal parts. Everyone will get a fraction of the whole quantity of sweets. One group is a quarter of the whole.
Half turn	A 180 degree rotation, i.e. $\frac{1}{2}$ of a 360 degree or 'full' turn.	
Hour	A unit of time.	There are 24 hours in one day.
Increasing	Becoming greater in value. Used in relation to number sequences.	2, 4, 6, 8. This number pattern is increasing by two each time.
Kilogram	A standard unit of mass, equal to 1000 grams.	The book has a mass of two kilograms .
Known fact	A number fact which has been committed to memory (or very fast recall) and can be applied fluently to various calculation strategies.	When I use the 'Make ten' strategy to add, I use known facts to partition the number I'm adding.
Left	Indicating the position or direction.	Make a quarter turn left and walk forward three steps.
Litre	A standard unit of volume, equal to 1000 millilitres.	The capacity of the jug is about half a litre .

Mental calculation	A calculation performed without using a formal written strategy. Simple jottings may aid a mental calculation.	14 plus 5 is equal to 19. I completed this using a mental calculation and deriving facts because I know that four plus five is equal to nine.
Metre	A standard unit of measure, equal to 100 centimetres.	I estimate that the table is about a metre tall.
Minute	A unit of time.	We will have lunch in five minutes .
Oblong	A quadrilateral with two pairs of parallel sides of equal length.	
Odd number	An integer which is not divisible by two without a remainder.	All numbers which end in 1, 3, 5, 7 and 9 are odd numbers .
Partition	To split a number into two or more parts.	The number 23 can be canonically partitioned (by place value) into 20 and 3, or non-canonically partitioned in many different ways, including 18 and 5, 17 and 6, etc.
Place value	A system for writing numbers, in which the value of a digit is defined by its position within the number.	In the number 452 written in base ten, the digit four has a value of 400, the five has a value of 50 and the two has a value of two.
Position	Location, expressed either descriptively using positional prepositions, or specified by coordinates.	The book is on the table. The clock is hanging above the board.
Pound (sterling)	The official currency of the United Kingdom.	Pounds sterling are written using the £ symbol. There are 100 pence in one pound sterling .
Property	Any attribute.	A property of a triangle is that it has three straight sides and three vertices, the sum of whose angles is 180 degrees.
Pyramid	A 3-D shape with a polygonal base and otherwise triangular faces, which form edges with the base, and which meet at an apex.	
Quantity	An amount, in some cases given a numerical value.	A quantity of apples is placed on the left-hand side of the balance. How many kilogram masses will we need to place on the right to balance the apples?
Quarter	One of four equal parts of a whole, quantity or object.	I have shared the eight conkers into four equal groups – I have two conkers, which is one quarter of the whole.
Quarter turn	A 90-degree rotation, i.e. $\frac{1}{4}$ of a 360 degree 'full' turn.	
Repeated addition	A structure of multiplication where equal parts are added to make a whole.	I can show 4×5 as repeated addition : $4 + 4 + 4 + 4 + 4$.



Mathematics Mastery

Repeated subtraction	A structure of division, where equal parts are subtracted and the number of equal parts summed to calculate a quotient.	I can use repeated subtraction to calculate 20 divided by four: $20 - 4 - 4 - 4 - 4 - 4$.
Represent	To express or show a mathematical concept using words, numerals and symbols, pictures, diagrams, or concrete manipulatives.	I have used three blue cubes to represent the three oranges in the question. I used a part-whole model to represent the addition question.
Right	Indicating the position or direction.	The picture is on the right -hand side of the board.
Rule	A consistent pattern which allows generalisation. Awareness of a rule allows a pupil to continue a sequence or generate a related sequence.	2, 5, 8, 11, 14... The rule is that each number is three greater than the previous number. Therefore, the next number in this sequence will be 17.
Scales	An object used to measure mass.	The scales showed that the banana had a greater mass than the apple.
Sign	Synonymous with symbol in its mathematical context, e.g. +, -, ×, ÷, =.	$20 \square 5 = 4$. What is the missing sign ?
Standard unit	A uniform measure, agreed upon as standard.	Standard units of mass include grams and kilograms. Standard units of length include centimetres, metres and kilometres. Standard units of volume and capacity include millilitres and litres.
Sphere	A 3-D shape with a continuous surface, which is at all points equidistant from its centre. It has an infinite number of flat faces and straight edges.	A bowling ball is a sphere .
Symbol	Synonymous with sign in its mathematical context, e.g. +, -, ×, ÷, =.	$20 \square 5 = 4$. What is the missing symbol ?
Table	A structure organised into columns and rows, in which data can be recorded.	The information for Thursday is not yet complete on the table because it is only Wednesday.
Turn	Rotation (see half and quarter turn).	A whole turn is 360 degrees. A half turn is 180 degrees. A quarter turn is 90 degrees.
Unit	<ol style="list-style-type: none"> 1. An element considered as a single entity. Ten single cubes can be grouped together to make a unit of ten. 2. A unit of measure, which can be standard or non-standard. 	I regrouped ten ones for one unit of ten. Unifix cubes can be used as units of measure, but these are not standard units .
Volume	A quantity or amount of any substance and the 3-D space it fills.	The bottle contains a volume of one litre but its capacity is two litres. The bottle is half full.



Mathematics Mastery

Year 2	Definition	Example
Angle	The amount of turn, measured in degrees.	The angle is 60 degrees.
Calculate	To compute or work out mathematically.	Can you calculate the answer to $13 + 4$?
Centimetre	A metric unit of length.	The book is 15 centimetres long.
Column	A vertical arrangement of numbers or objects.	23 has two tens – I will place them into the tens column .
Commutative	A property of addition and multiplication. It does not matter in which order the addends or factors are added or multiplied; the result will be the same.	$4 + 6 = 10$ $6 + 4 = 10$ This demonstrates that addition is commutative . Arrays demonstrate the commutativity of multiplication, i.e. $3 \times 4 = 4 \times 3$
Consecutive	Following in order.	2, 3, 4, 5, 6 are consecutive numbers. 3, 6 and 9 are consecutive multiples of 3.
Denominator	The number written below the vinculum in a fraction. In a measure context, it indicates the number of equal parts into which the whole is divided. In a division context, it is the divisor.	In the fraction one quarter, four is the denominator .
Division	The process of partitioning a whole into equal parts.	12 divided by 3 is equal to 4.
Efficient	Well-organised. Choosing an efficient computation strategy requires consideration of the numbers involved and will normally utilise 'known facts'.	I will use my number bonds knowledge to calculate $22 + 7$ efficiently . I know that $2 + 7$ is equal to 9, so the answer is 29. That's more efficient than counting on seven.
Frequency	The number of times something occurs within a data set.	4 pupils have brown hair. The frequency of brown hair is 4.
Gram	A metric unit of mass.	The pencil weighs 20 grams .
Heptagon	A polygon with seven sides and seven angles.	
Hexagon	A polygon with six sides and six angles.	
Inverse operations	Opposite operations that 'undo' each other.	Addition and subtraction are inverse operations.
Millilitre	A metric unit of capacity/volume.	The can of fizzy drink has a capacity of 330 millilitres .
Multiple	The result of multiplying a number by an integer, for example, 12 is a multiple of 3 and 4 because $3 \times 4 = 12$.	36 is a multiple of three because three multiplied by 12 is equal to 36. It is also a multiple of 12 for the same reason (and 1, 2, 4, 6, 9, 18 and 36).
Multiplication	One of the four mathematical operations. Multiplication can be understood as repeated addition or scaling (introduced in Year 3).	The multiplication symbol is \times .
Multiply	To increase a quantity by a given scale factor.	I can multiply 3 by 4 which is equal to 12.



Near double	When two numbers involved in an addition are close in value, such as $23 + 22$. The numbers can be treated as exact doubles, followed by compensating.	To calculate $23 + 22$, I can use the near double strategy. I can double 22 and then add one more.										
Non-unit fraction	A fraction with a numerator greater than one.	Two thirds is a non-unit fraction .										
Numerator	The number written above the vinculum in a fraction. In a measure context, it indicates the specified number of parts out of the whole. In a division context, it is the dividend.	In the fraction one quarter, one is the numerator .										
Octagon	A polygon with eight sides and eight angles.											
Operation	A mathematical process. The four mathematical operations are addition, subtraction, multiplication and division.	$4 + 2 = 6$. The operation is addition.										
Pentagon	A polygon with five sides and five angles.											
Pictogram	A representation of data using pictures or symbols.	<p>Countries people visited</p> <table border="1"> <tr> <td>France</td> <td>●●●●</td> </tr> <tr> <td>Germany</td> <td>●●●●●</td> </tr> <tr> <td>America</td> <td>●●●</td> </tr> <tr> <td>China</td> <td>●</td> </tr> <tr> <td>Australia</td> <td>●</td> </tr> </table> <p>Each ● stands for 10 people</p>	France	●●●●	Germany	●●●●●	America	●●●	China	●	Australia	●
France	●●●●											
Germany	●●●●●											
America	●●●											
China	●											
Australia	●											
Quadrilateral	A 2D shape with four sides and four angles. which add up to 360 degrees.											
Relationship	The way in which two or more things are connected.	The relationship between addition and subtraction is that they are the inverse of each other.										
Right angle	An angle of 90 degrees.	A square has four right angles .										
Rotation	The act of rotating about an axis/centre.	I will rotate the square 90 degrees clockwise.										
Scale	Equally spaced markings on a measuring device which can be read to quantify a measurement.	Using the scale on the ruler, the book measures 15cm.										
Symmetry	A shape is symmetrical when it fits exactly onto itself when folded in half.	This triangle has one line of symmetry . 										
Tally	A form of counting. Each tally is a vertical mark. After the fourth vertical mark, a fifth	Four children have black hair; I will record this as four tallies .										





	horizontal/diagonal mark is drawn to create a group of five.	
Temperature	The measure of heat.	Outside has a temperature of 15 degrees Celsius.
Unit fraction	A fraction with a numerator of one.	One-third is a unit fraction .
Vinculum	A horizontal line that separates the numerator and the denominator in a fraction.	$\frac{1}{4}$ ← vinculum


Year 3	Definition	Example
Acute angle	An angle that is smaller than a right angle.	It is smaller than my right angle checker so this must be an acute angle .
Axis (plural: axes)	A real or imaginary reference line. The y-axis (vertical) and x-axis (horizontal) on charts and graphs are used to show the measuring scale or labels for the variables.	The y-axis on this bar graph shows you how many pupils preferred each colour.
Bar graph	A representation of data in which the frequencies are represented by the height or length of the bars.	This bar graph shows us the preferred colours of the pupils in our Year 3 class.
Columnar addition/subtraction	The formal written algorithms for addition and subtraction that are exemplified in <i>Mathematics Appendix 1</i> of the 2014 national curriculum.	Solve the following calculations by using the appropriate method of columnar addition or subtraction .
Factor	A number, that when multiplied with one or more other factors, makes a given number.	The number six has four factors: 1, 2, 3 and 6.
Formal written methods	Exemplified in <i>Mathematics Appendix 1</i> (see above). As well as including columnar addition and subtraction, these also consist of written algorithms for multiplication and division.	Pupils should only use formal written methods for calculations that cannot be efficiently calculated using mental strategies (with or without jottings).
Horizontal	Horizontal refers to planes and line segments that are parallel to the horizon.	The x-axis on a graph should be horizontal .
Irregular	In geometry, irregular is a term used to describe shapes that are not regular (see below).	The sides and the angles of this pentagon are not all equal so the pentagon is irregular . 
Kilometre	A metric unit measure of length that is equal to one thousand metres.	The distance from the school to Arun's house was exactly one kilometre .
Millimetre	A metric unit measure of length that is equal to one thousandth of one metre.	The length of Philippa's ruler is 300 millimetres .



Numeral	A numeral is a symbol (or group of symbols) used to represent a number.	Whole numbers can all be represented as numerals consisting of the digits 0 to 9.
Obtuse angle	An angle that is greater than a right angle but less than 180 degrees.	It is greater than my right angle checker so this angle must be obtuse.
Parallel	Line segments that can be described as parallel must be on the same plane and will never meet, regardless of how far either or both line segments are extended.	The opposite sides of a square are parallel .
Perimeter	The perimeter of a 2-D shape is the total distance around its exterior.	I know that one side of this square is 2cm so it must have a perimeter of 8cm.
Perpendicular	A pair of line segments (or surfaces) can be described as perpendicular if they intersect at (or form) a right angle.	The adjacent sides of a rectangle are perpendicular .
Place holder	A place holder is a zero used in any place value column (that contains a value of zero) to clarify the relative positions of the digits in other places.	I need to use a place holder in the ones column to make it clear that my number is 320 and not 32.
Prism	A prism is a 3-D solid with two identical, parallel bases and otherwise rectangular faces.	A triangular prism has five faces, consisting of three rectangles and two triangles which are parallel.
Product	The result you get when you multiply two numbers.	24 is the product of 3 and 8.
Regular	Regular 2-D shapes (regular polygons) have angles that are all equal and side lengths that are all equal. Regular 3-D shapes (the Platonic Solids) are those that have congruent (exactly the same) faces of a single regular polygon.	A square is a regular 2-D shape because all four angles are right angles and all four sides are the same length. A cube is a regular 3-D shape with six identical square faces.
Roman numeral	Roman numerals are a system of symbols used to represent numbers that were developed and used by the Romans. They do not use a place value system.	The number twelve on this clock is represented by the Roman numerals XII, which is $10 + 1 + 1$.
Round	Approximate a number, normally to the nearest multiple of ten, to make it easier with which to calculate.	I would round the number 17 to 20 because it is three away from 20 but seven away from 10.
Square-based pyramid	A pyramid is a 3-D shape with a 2-D shape (which gives the pyramid its name) as a base and triangular faces that taper to a point called a vertex or apex.	This square-based pyramid has five faces; one square face and four triangular faces.
Triangle-based pyramid		This triangle-based pyramid has four triangular faces.


Year 4	Definition	Example
Area	The space a surface takes up inside its perimeter. Area is always measured in square units.	 <p>The area is 8 square units.</p>
Associative law	No matter how the parts in an addition or multiplication equation are grouped, the answer will be the same.	$(6 + 3) + 2 = 11$ $6 + (3 + 2) = 11$ Addition and multiplication are associative. Subtraction and division are not.
Convert	To change from one unit of measurement to another.	2 km can be converted to metres – it is equal to 2000 m.
Coordinate	The position of a point, usually described using pairs of numbers. Sometimes called Cartesian coordinates, after the mathematician Rene Descartes.	The coordinate (3,4) describes a point that is 3 on the x axis and 4 on the y axis.
Decimal fraction	A fraction expressed in its decimal form.	Half written as a decimal fraction is 0.5.
Distributive law	The process whereby adding some numbers and then multiplying the sum gives the same answer as multiplying the numbers separately and then adding the products.	$3 \times (2 + 4) = (3 \times 2) + (3 \times 4)$ $3 \times 12 = (3 \times 10) + (3 \times 2)$
Dividend	The amount that you want to divide.	In ' $12 \div 3 = 4$ ', 12 is the dividend.
Divisor	The number you divide by.	In ' $12 \div 3 = 4$ ', 3 is the divisor.
Equilateral	Having all sides the same length.	An equilateral triangle has three equal sides.
Equivalent	Equivalent means having the same value. Equivalent fractions have the same value.	$\frac{2}{4} = \frac{1}{2}$
Expression	One or a group of numbers, symbols or operators. An expression does not use equality or inequality signs. <i>Using an equality or inequality sign will give an equation.</i>	2×3 4^2
Grid	A series of evenly divided and equally spaced shapes, usually squares.	
Improper fraction	A fraction where the numerator is bigger than the denominator. These fractions are therefore greater than one whole.	$\frac{12}{11}$
Integer	A whole number that can be positive or negative.	6 is an integer, 0.6 is not.
Interval	An interval on a graph's axis lies between two values.	If one point on an axis is 50 and the next 60, the interval is 10.





Isosceles	Having two sides of equal length. Isosceles triangles have two equal sides; isosceles trapezia have two equal, non-parallel sides.	
Kite	A 2-D shape with two pairs of equal length adjacent sides. The diagonals intersect at right angles.	
Line graph	A graph that uses lines to connect the points on a data chart. Used to present continuous data, such as change over time.	
Mixed numbers	Numbers consisting of an integer and fractional part.	$1\frac{1}{2}$, $3\frac{3}{4}$
Negative number	A number that is less than zero. (It is helpful to refer to these numbers as 'negative numbers' rather than 'minus' to avoid confusion with the operation 'minus'.)	-1, -24, -0.5 etc.
Parallelogram	A 2-D shape that has two pairs of parallel sides and equal opposite angles.	
Plot	To mark out a point on a graph or grid.	'Plot the point (3,6)' means to draw the precise location of that point, usually shown as a dot or a small cross.
Point	The precise location of a position on a 2-D plane.	An exact place on a graph or on squared paper. A point is often represented by a capital letter. 
Positive number	A number that is greater than zero. Zero is neither positive or negative.	3, 32, 0.5
Proper fraction	A fraction with a value less than one.	$\frac{1}{2}$, $\frac{3}{4}$, $\frac{5}{8}$
Protractor	A measuring device for measuring the size of an angle. Angles are measured in degrees (°).	
Quotient	The result when the dividend is divided by the divisor.	$15 \div 3 = 5$ 5 is the quotient .
Rectilinear	A rectilinear shape has straight line edges which are perpendicular (all meet at right angles).	A rectangle.  A straight-sided shape that can be divided up into other rectangles. 





Rhombus	An equilateral parallelogram with four equal length sides.	
Scalene	A scalene triangle has three unequal sides and three unequal angles.	
Short division	A formal written layout where the quotient is calculated showing only one written step.	$\begin{array}{r} 77 \\ 5 \overline{) 385} \end{array}$
Short multiplication	A formal written layout where the multiplier is usually 9 or less.	$\begin{array}{r} 782 \\ \times 9 \\ \hline 7038 \end{array}$
Simplify	To write a number or equation in its simplest form.	I can simplify $\frac{8}{10}$ to $\frac{4}{5}$.
Square centimetre	A unit of measure for area equal to a square with the dimensions 1 cm by 1cm.	Sometimes referred to as centimetre squared, abbreviated to cm^2 .
Trapezium	A quadrilateral with exactly one pair of parallel sides.	

Year 5	Definition	Example
Angle at a point	Angles that meet at a point that sum to 360° .	
Angle on a line	Angles formed on a straight line that sum to 180° .	
Average (mean)	A measure of central tendency. The mean average of a set of data is the sum of the quantities divided by the number of quantities.	The mean average of the set 4, 5, 5, 6 is 5 because $(4 + 5 + 5 + 6) \div 4 = 5$.
Common factor	A factor of two (or more) given numbers.	A common factor of 12 and 9 is 3 because $3 \times 4 = 12$ and $3 \times 3 = 12$.
Common multiple	A multiple of two (or more) given numbers.	A common multiple of 3 and 6 is 12 because $3 \times 4 = 12$ and $6 \times 2 = 12$.
Congruent	Used to describe two shapes or figures which are exactly the same size.	The two triangles are congruent. If I place one on top of the other, there is no overlap. 
Cube number	The product of three equal factors.	Eight is a cube number because $8 = 2 \times 2 \times 2 = 2^3$.
Cubic centimetre	A unit used to measure volume. The space taken up by a cube with edges of length 1 cm or	The volume of this multilink cube is eight cubic centimetres.





	which measures $1\text{ cm} \times 1\text{ cm} \times 1\text{ cm}$.	
Cubic metre	A unit used to measure volume. The space taken up by a cube with edges of length 1 metre.	The volume of this fridge is two cubic metres.
Decagon	A polygon with ten sides and ten angles.	
Degree	The unit of measure for angles.	A right angle is 90 degrees.
Diagonal	A straight line segment that joins one vertex to another.	The diagonals of a kite are perpendicular 
Divisible	A number is said to be divisible by another if it can be divided by that number without a remainder.	24 is divisible by 8. When divided by 8 it gives a quotient of 3, with no remainder.
Dodecagon	A polygon with twelve sides and twelve angles.	
Long division	The formal written algorithm that can be used to divide by a number with two or more digits.	$\begin{array}{r} 34 \\ 12 \overline{) 408} \\ \underline{36} \\ 48 \\ \underline{48} \\ 0 \end{array}$
Long multiplication	The formal written algorithm that can be used to multiply a number by a number with two or more digits.	$\begin{array}{r} 34 \\ \times 12 \\ \hline 68 \\ 340 \\ \hline 408 \end{array}$
Negative integer	A whole number with a value less than zero. Zero is neither positive nor negative.	When the temperature falls below 0° a negative integer is used to record it.
Nonagon	A polygon with nine sides and nine angles.	
Percentage	The number of parts per hundred which is written using the % symbol.	30% means for every 100 there are 30.
Polygon	A 2-D shape with three or more straight sides.	Triangles and rectangles are examples of polygons.
Polyhedron (pl. polyhedra)	A 3-D shape with flat surfaces that are polygons.	A cuboid is a polyhedron. A cylinder is not a polyhedron because it has a curved surface.
Prime factor	A factor that is a prime number.	3 and 2 are prime factors of 6.


Prime number	A whole number with only two factors, one and the number itself.	2, 3, 5, 7, 11, 13, 17 and 19 are the prime numbers less than 20.
Remainder	The amount remaining after division when a whole number answer is needed.	21 divided by four is equal to five with a remainder of 1.
Reflection	A mirror image that is equidistant from a mirror line.	The shape has been reflected in the dotted mirror line. 
Reflex angle	An angle that is greater than 180°.	
Scale (not to scale)	The ratio of lengths, in a drawing, are in proportion to the measurements of the real object. The lengths are not in proportion when not to scale.	The diagram was not drawn to scale. That means I can't use a ruler to measure the sides, because they are not in proportion to the real object.
Square metre	A unit of measure for area. The surface covered by a square with sides of length one metre.	The area of the floor in a room might be measured in square metres .
Square number	The product of two equal factors.	9 is a square number because $9 = 3 \times 3 = 3^2$.
Tetrahedron	A 3-D shape with four triangular faces.	
Transformation	A collective term for the ways that shapes can be changed, resulting in congruent or similar shapes, i.e. translation, reflection, rotation or enlargement.	Translations and reflections are types of transformations .
Translation	When a shape moves so that it is in a different position but retains the same size, area, angles and side length and so is congruent.	Triangle C has been translated three right and two down resulting in triangle D. 

Year 6	Definition	Example
Arc	A portion of the circumference of a circle	
Brackets	The symbols () used to separate parts of a multi-step calculation.	$(10 - 2) \times 3 = 24$
Centre	In a circle, the centre refers to one point that is equidistant to all points around the circumference of the circle.	To draw a circle, I place the point of my pair of compasses at the centre.
Circumference	The perimeter/boundary of a circle.	
Compasses	A tool for creating curved lines, arcs and circles.	I can use a pair of compasses to draw a circle with a radius of 4 cm.
Common fraction	A fraction written with a numerator and denominator separated by a vinculum.	One quarter can be written as a common fraction, $\frac{1}{4}$.
Degree of accuracy	A description of how accurately a value is communicated.	The degree of accuracy needed for the answer is one decimal place.
Diameter	A line from one point of the circumference of a circle to another on the opposite side, which must pass through the centre of the circle.	The circumference of a circle is the diameter multiplied by pi. 
Equivalent expression	An expression, which can be algebraic, which is equal in value to another expression.	Find an equivalent expression to $17 + 10$. $18 + 9$ is an equivalent expression to $17 + 10$.
Factorise	To identify factors of a given number. To express a number as factors.	I can factorise 12 by looking at its factor pairs. $1 \times 12 = 12$, $2 \times 6 = 12$, $3 \times 4 = 12$. So the factors of 12 are 1, 2, 3, 4, 6 and 12.
Foot/feet	An imperial unit of measure of length.	I am approximately five feet tall.
Formula	An algebraic expression of a rule.	The area of a rectangle can be found by multiplying the width and height. $a = w \times h$
Gallon	An imperial unit of measure of volume/capacity.	A gallon is approximately 4.5 litres
Imperial unit	A unit of measure once officially used in the UK but is now used less often, except in the context of length. Includes miles, pounds and pints.	Miles are an imperial unit to measure length.
Inch	An imperial unit of measure.	An inch is approximately 2.2 cm.
Intersect	The point at which two (or more) lines meet is where they intersect.	The x and y axes intersect at (0,0)



Metric unit	A standard unit of measure used in the UK and Europe. Includes centimetres, litres and kilograms.	Litres are a metric unit used to measure volume.
Mile	An imperial unit of measure of length.	Five miles is equivalent to eight kilometres.
Net	A group of 2-D shapes which, when folded and connected, forms a 3-D polyhedron.	The net of a cube is comprised of six connected squares.
Order of operations	The internationally agreed order to complete operations in a multi-step equation with multiple operations.	$(3 + 4) \times 2 = \square$ The order of operations dictates that the operation within the brackets is completed first.
Origin	The point at which axes in a coordinates grid cross; the point (0,0).	 <p>The origin is indicated by the blue dot.</p>
Ounce	An imperial unit of measure of mass.	The newborn baby had a mass of 6 pounds and 3 ounces.
Pie chart	A representation of a set of data where each segment represents one group in proportion to the whole.	<p>Nationality of Astronauts on Board ISS January 2017</p>  <p>■ Russia ■ USA ■ France</p>
Pint	An imperial unit of measure.	I found a pint of milk on my doorstep.
Pound (mass)	An imperial unit of measure of mass.	The new-born baby had a mass of 6 pounds and 3 ounces.
Proportion	A comparison between two or more parts of a whole or group. Proportion expresses a part-whole relationship. This may be represented as a fraction, a percentage or a decimal.	Two thirds of a class were boys. The proportion of the class that is girls is one third.
Quadrant	One of four regions into which a coordinates grid is divided.	



Radius	A line from one point of the circumference of a circle to the centre of the circle.	
Ratio	A comparison between two or more parts of a whole or group. Ratio expresses a part-part relationship. This is usually represented in the form a:b.	For every 4 tulips there are 7 daffodils. The ratio of tulips to daffodils is 4:7.
Similar	Similar shapes are those which have the same internal angles and where the side lengths are in the same ratio or proportion. Enlarging a shape by a scale factor (for example by doubling all side lengths) creates a similar shape.	All squares are similar to one another.
Square millimetre	The area of a square with sides 1 mm.	The smallest squares on graph paper have an area of one square millimetre.
Square kilometre	The area of a square with sides 1 km.	The area of England is 130 279 square kilometres.
Vertically opposite angles	Angles which are positioned opposite to one another when two lines intersect.	 The purple angles indicated are vertically opposite angles.